CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Level

MARK SCHEME for the May/June 2014 series

9231 FURTHER MATHEMATICS

9231/22 Paper 2, maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2014 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

CAMBRIDGE
International Examinations

Question Number	Mark Scheme Details		Part Mark	Total
1	Equate impulse to momentum to find initial speed v and Newton's law of restitution to find new speed:	$v = 4u,\ v' = ev = [-]\,3u$ M1 A1	2	**2**
2	Find v^2 at both A and B:	$v_A^{\,2} = \omega^2(a^2 - 0{\cdot}5^2)$ *and* $v_B^{\,2} = \omega^2(a^2 - 0{\cdot}75^2)$ B1		
	Find amplitude a m from given K.E. ratio:	$\frac{1}{2} mv_A^{\,2} = (12/11)\ \frac{1}{2} mv_B^{\,2}$		
		$11\,(a^2 - 0{\cdot}5^2) = 12\,(a^2 - 0{\cdot}75^2)$		
		$a^2 = \frac{1}{4}(27 - 11) = 4,\ \ a = 2$ M1 A1	3	
	Find ω from $v_{\max} = a\omega$:	$0{\cdot}6 = 2\omega,\ \ \omega = 0{\cdot}3$ B1		
	Find time ($\surd$ on a) at A	$\omega^{-1}\sin^{-1}(0{\cdot}5/2)$ *or* $\omega^{-1}\cos^{-1}(0{\cdot}5/2)$		
	or at B, e.g.:	$\omega^{-1}\sin^{-1}(0{\cdot}75/2)$ *or* $\omega^{-1}\cos^{-1}(0{\cdot}75/2)$ M1 A1$\surd$		
	Combine correctly to find time from A to B:	$\omega^{-1}\sin^{-1}(0{\cdot}75/2) - \omega^{-1}\sin^{-1}(0{\cdot}5/2)$		
		or $\omega^{-1}\cos^{-1}(0{\cdot}5/2) - \omega^{-1}\cos^{-1}(0{\cdot}75/2)$ M1		
	Evaluate to 3 d.p.:	$= \omega^{-1}(0{\cdot}3844 - 0{\cdot}2527)$ or $\omega^{-1}(1{\cdot}318 - 1{\cdot}186)$		
		$= 1{\cdot}2813 - 0{\cdot}8423$		
		$4{\cdot}3937 - 3{\cdot}9547 = 0{\cdot}439$ [s] A1	5	**8**

© Cambridge International Examinations 2014

3	Use conservation of momentum, e.g.:	$mv_A + 9mv_B = mu$	M1		
	Use Newton's law of restitution (consistent signs):	$v_B - v_A = eu$	M1		
	Relate v_A to v_B using K.E. (A.E.F.):	$\frac{1}{2}mv_A^2 + \frac{1}{2}9mv_B^2 = \frac{1}{4}mu^2$	M1		
	Combine two eqns to find v_A and v_B e.g.:	$v_A = (1 - 9e)u/10$, $v_B = (1 + e)u/10$			
		or v_A , $v_B = -u/2$, u/6 [or $7u/10$, $u/30$]	M1 A1		
	Use in 3rd eqn to find e, e.g.:	$(1 - 9e)^2 + 9(1 + e)^2 = 50$			
	(A0 if finally $\pm\frac{2}{3}$)	$90e^2 = 40$, $e = \frac{2}{3}$	M1 A1	7	
	Use Newton's law of restitution with	$v_C = 2v_B'$, e.g.: $v_C - v_B' = ev_B$, $v_B' = \frac{2}{3}v_B$	B1		
		[$v_B = u/6$, $v_B = u/9$, $v_C = 2u/9$]			
	Use conservation of momentum to find k:	$9mv_B' + kmv_C = 9mv_B$			
		$9v_B' + 2kv_B' = 13{\cdot}5v_B'$, $k = 9/4$	M1 A1	3	**10**

© Cambridge International Examinations 2014

4	**(i)**	Use conservation of energy at lowest point: Use $F = ma$ radially at lowest point: Eliminate v^2 to find R $[v^2 = 2{\cdot}3\,ga]$:	$\frac{1}{2}mv^2 = \frac{1}{2}mu^2 + mga$ $R - mg = mv^2/a$ $R = mu^2/a + 3mg = 3{\cdot}3\,mg$	B1 B1 B1	3	
	(ii)	Use conservation of energy at B to find V_B : (A.E.F.)	$\frac{1}{2}mV_B^2 = \frac{1}{2}mu^2 + mga\sin\theta$ $V_B^2 = (0{\cdot}3 + 0{\cdot}5)ga$, $V_B = \sqrt{(0{\cdot}8\,ga)}$ *or* $2\sqrt{(ga/5)}$ *or* $0{\cdot}894\sqrt{(ga)}$	M1A1 A1	3	
	(iii)	Use vertical component v_B of speed V_B at B: Find height h reached above B: Find height h reached above level of O:	$v_B = V_B\cos\theta$ $[= \frac{1}{4}\sqrt{15}\,V_B = \sqrt{(\frac{3}{4}ga)}]$ $h = v_B^2/2g = 3a/8$ $h - a\sin\theta = 3a/8 - \frac{1}{4}a = a/8$ **A.G.**	M1 M1 A1 A1	4	**10**
5		Find MI of components about A: (M1 for BC or CD) Find total MI about A: (OR can first find total MI about centre of mass) State or imply total mass acts at mid-point of AC Use eqn of circular motion to find $d^2\theta/dt^2$: Approximate $\sin\theta$ by θ and substitute for I: Find period $T = 2\pi/\omega$ with $\omega = \sqrt{(49\,g/384\,a)}$:	Glass $(3M/5)\{\frac{1}{3}(5a)^2 + 25a^2\} = 20\,Ma^2$ AB $M\{\frac{1}{3}(4a)^2 + (4a)^2\} = 64\,Ma^2/3$ AD $\frac{1}{3}M\{\frac{1}{3}(3a)^2 + (3a)^2\} = 4\,Ma^2$ BC $\frac{1}{3}M\{\frac{1}{3}(3a)^2 + 73a^2\} = 76\,Ma^{2/3}$ CD $M\{\frac{1}{3}(4a)^2 + 52a^2\} = 172\,Ma^{2/3}$ $I = 128\,Ma^2$ A.G. $I\,d^2\theta/dt^2 = [-]\,(49Mg/15)\,5a\sin\theta$ $d^2\theta/dt^2 = -(49g/384a)\,\theta$ $T = 2\pi\sqrt{(384a/49g)}$ *or* $(16\pi/7)\sqrt{(6a/g)}$ *or* $17{\cdot}6\sqrt{(a/g)}$ (A.E.F.)	M1 A1 B1 B1 M1 A1 A1 A1 M1 M1 A1 A1 B1	8 5	**13**

© Cambridge International Examinations 2014

6		State or find the expected value of X: using $p = ¼$:	$E(X) = 1/p = 1/¼ = 4$ B1	1	
	(i)	Find $P(X = 4)$:	$P(X = 4) = (¾)^3 ¼ = 27/256$ or 0·105 M1 A1	2	
	(ii)	Find $P(X<6)$:	$P(X < 6) = 1 – (¾)^5$ or $\{1 + ¾ + (¾)^2 + (¾)^3 + (¾)^4\}¼$		
		S.R. Using $p = ½$ can earn B0 M1 A0 M0 A0	$= 781/1024$ or 0·763 M1 A1	2	**5**
7	**(i)**	State probability density function of T:	$f(t) = 0·001 \exp(-0·001t)$ $(t \geq 0)$ B1 [= 0 (otherwise *or* $t < 0$)]	1	
	(ii)	Find $P(T > 2000)$:	$P(t > 2000) = 1 – F(2000)$ M1		
		S.R. $1 – e^{-2} = 0·865$ earns B1 only (max 1/3)	$= 1 – (1 – e^{-2}) = e^{-2}$ *or* 0·135 M1 A1	3	
		State inequality for t (lose A1 if = or ≤):	$(\exp(-0·001t))^{10} \geq$ [*or* >] 0·9 M1 A1		
		Solve for t_{max}:	$t_{max} = (\ln 0·9)/(-0·01) = 10·5$ M1 A1	4	**8**
		(Omitting power 10 earns 0/4; using $1 – (\exp(-0·001t))^{10}$ can earn M1 A0 M1 A0 only)			

© Cambridge International Examinations 2014

8	State hypotheses (B0 for $\bar{\chi}$ …):	$H_0: \mu_X = \mu_Y$, $H_1: \mu_X \neq \mu_Y$	B1		
	Estimate both popln. variances using two samples: (allow use of biased: $\sigma_{X,60}^2 = 236$ *or* $15{\cdot}36^2$)	$S_x^2 = (626\,220 - 6060^2/60) / 59$ [= 240 *or* $15{\cdot}49^2$]			
	(allow use of biased: $\sigma_{Y,50}^2 = 265$ *or* $16{\cdot}28^2$)	*And* $s_Y^2 = (464\,500 - 4750^2/50) / 49$ [= 270·4 *or* $16{\cdot}44^2$]	M1 A1		
	Estimate population variance for combined sample: (allow $\sigma_{X,60}^2/60 + \sigma_{Y,50}^2/50$: 9·233 *or* $3{\cdot}039^2$)	$s^2 = s_X^2/60 + s_Y^2/50$ $= 9{\cdot}408$ *or* $3{\cdot}067^2$	M1 A1		
	Calculate value of z (to 2 d.p., either sign):	$z = (101 - 95) / s$	M1		
		$= 6/3{\cdot}067 = 1{\cdot}96$ (*or* 1·97)	A1		
	State or use correct tabular z – value (to 2 d.p.): (or can compare 6 with e.g. 2·326 s = 7·13 *or* 7·07)	$z_{0.99} = 2{\cdot}326$ *or* 2·33 (allow 2·36)	B1		
	Correct conclusion (A.E.F, √on z – values):	[Accept H_0] Claims are the same	B1√		
	S.R. Assuming equal population variances:	Hypotheses; Explicit assumption	(B1; B1)		
	Find pooled estimate of common variance s^2:	: $s^2 = (626\,220 - 6060^2/60 +$ 464 500 – $4750^2/50) / 108$	(M1 A1)		
	Calculate value of z (to 2 d.p., either sign):	$z = 6 / s\sqrt{(1/60+1/50)} = 1{\cdot}97$	(M1 A1)		
		$= 253{\cdot}8$ *or* $15{\cdot}93^2$	(A1)		
	Tabular value; conclusion	As above)	(B1; B1√		**9**

© Cambridge International Examinations 2014

9	Find expected frequency p:	$p = 200 \int_2^3 (1 / x \ln 8)\, dx$			
		$= (200 / \ln 8) [\ln x]_2^3$			
		$= 200 \times 0{\cdot}1950 = 39{\cdot}00$ **A.G.**	M1A1		
		$q = 21{\cdot}46$ *or* $21{\cdot}45$	M1A1		
	Find q by similar method *or* by using total of 200:			4	
	State (at least) null hypothesis:	H_0: f (x) fits data (A.E.F.)	B1		
	Calculate χ^2 (to 3 s.f.):	$\chi^2 = 0{\cdot}202 + 0{\cdot}923 + 0{\cdot}678 + 0{\cdot}584$			
		$+ 1{\cdot}134 + 4{\cdot}134 + 3{\cdot}644 = 11{\cdot}3$	M1A1		
	State or use correct tabular χ^2 value (to 3 s.f.)::	$\chi^2_{6,\,0.95} = 12{\cdot}59$	B1		
	Valid method for reaching conclusion:	Accept H_0 if $\chi^2 \leqslant$ tabular value	M1		
	Conclusion consistent with correct values (A.E.F):	Distribution fits observations	A1	6	**10**
10	Find correlation coefficient r:				
	$r = (73\,527 - 866 \times 639 / 10) / \sqrt{\{(121\,276 - 866^2 / 10)(55\,991 - 639^2 / 10)\}}$		M1 A1		
	(A.E.F.; A0 if only 3 s.f. clearly used)	$= 18\,189{\cdot}6 / \sqrt{(46\,280{\cdot}4 \times 15\,158{\cdot}9)}$	A1		
		$= 0{\cdot}687$	*A1	4	
	State both hypotheses (B0 for r …):	H_0: $\rho = 0$, H_1: $\rho \neq 0$	B1		
	State or use correct tabular two-tail r-value:	$r_{10,\,5\%} = 0{\cdot}632$	*B1		
	Valid method for reaching conclusion:	Reject H_0 if $\lvert r \rvert >$ tabular value	M1		
	Correct conclusion (A.E.F, dep *A1, *B1):	There is non-zero correlation	A1		
	Calculate gradient p in $x - \bar{\chi} = p(y - \bar{\gamma})$:	$p = 18\,189{\cdot}6 / 15\,158{\cdot}9 = 1{\cdot}20$	B1	4	
	Find regression line of x on y:	$x = 86{\cdot}6 + 1{\cdot}20(y - 63{\cdot}9)$			
		$= 1{\cdot}20\,y + 9{\cdot}92$	M1 A1	3	**11**

© Cambridge International Examinations 2014

11 A (i)	Use Pythagoras to find AB:	$AB = \sqrt{(4a^2 + 12a^2)} = 4a$	**A.G.**	M1 A1		
	Find $\angle SAB$:	$\angle CAB = \sin^{-1} 2a\sqrt{3}/4a$ *or* $\cos^{-1} 2a/4a$				
		or $\tan^{-1} 2a\sqrt{3}/2a$				
(ii)		$= 60°$ so $\angle SAB = 30°$	**A.G.**	M1 A1	4	
	EITHER					
	Resolve vertically and horizontally, e.g.:	$\frac{1}{2} N_A + \frac{1}{2}\sqrt{3}\, N_B + \frac{1}{2}\sqrt{3}\, F_A = W$				
	(F_A may be in either direction)	*and* $\frac{1}{2}\sqrt{3}\, N_A = \frac{1}{2} N_B + \frac{1}{2} F_A$		M1 A1		
	Eliminate $N_B + F_A$ to find N_A:	$N_A = \frac{1}{2} W$	**A.G.**	A1		
	OR				3	
(iii)						
	Resolve in dirn. PQ to find N_A:	$N_A = \frac{1}{2} W$	**A.G.**	(M1 A1)		
	Second resolution, e.g. in dirn. PS:	$N_B + F_A = \frac{1}{2}\sqrt{3}\, W$		(A1)		
	Take moments, e.g. about A:	$\frac{1}{2}\sqrt{3}\, W \times 3a/2 + \frac{1}{2} W \times (2\sqrt{3} - 3)a$				
	(A1 for each side of eqn)	$= N_B \times 2a$		M1 A1 A1		
	Solve to find N_B:	$N_B = \{(7\sqrt{3} - 6)/8\}\, W$		M1 A1		
	Use N_B to find F_A:	$F_A = \sqrt{3}\, N_A - N_B$ *or* $\frac{1}{2}\sqrt{3}\, W - N_B$				
		$= \{3(2 - \sqrt{3})/8\} W$ (A.E.F.)		M1 A1	7	**14**

© Cambridge International Examinations 2014

B	Estimate population variance:	$s_P^2 = (236{\cdot}0 - 42{\cdot}8^2/8)/7$			
	(allow biased here: $0{\cdot}8775$ or $0{\cdot}9367^2$)	$= 351/350$ or $1{\cdot}003$ or $1{\cdot}001^2$	M1		
	Find confidence interval (allow z in place of t) e.g.:	$42{\cdot}8/8 \pm t\sqrt{(s_P^2/8)}$	M1		
	Use correct tabular t-value:	$t_{7,\,0.975} = 2{\cdot}365$	A1		
	Evaluate C.I. correct to 2 d.p.:	$5{\cdot}35 \pm 0{\cdot}84$ or $[4{\cdot}51, 6{\cdot}19]$	A1	4	
	Formulate inequality for k (or equality for $k_{\max}$):	$(5{\cdot}35 - k)/\sqrt{(s_P^2/8)} \geqslant$ [or $>$] t	M1		
	Use correct tabular t-value:	$t_{7,\,0.9} = 1{\cdot}415$	A1		
	Solve for $k_{\max}$ (A0 if = or $\leqslant$ was used for k above):	$5{\cdot}35 - k \geqslant 0{\cdot}50$, $k_{\max} = 4{\cdot}85$	A1	3	
	State hypotheses (B0 for $\bar{x}$ …), e.g.:	H_0: $\mu_P = \mu_Q$, H_1: $\mu_P > \mu_Q$	B1		
	State assumption (A.E.F.):	Normal distns. for [P and] Q			
		and equal variances	B1		
	Estimate (pooled) common variance:	$s^2 = (7 \times 1{\cdot}003 + 11 \times 1{\cdot}962)/18$			
		$= 1{\cdot}589$ or $1{\cdot}261^2$	M1 A1		
	Calculate value of t (to 3 s.f.):	$t = (5{\cdot}35 - 4{\cdot}60)/(s\sqrt{(1/8 + 1/12)})$			
		$= 1{\cdot}30$	M1 A1		
	Correct conclusion (A.E.F., $\surd$ on t):	$t < t_{18,\,0.9} = 1{\cdot}33$ so Q's mean is not less than P's	B1$\surd$	7	**14**

© Cambridge International Examinations 2014