Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

2380869687

FURTHER MATHEMATICS **9231/13**

Paper 1 Further Pure Mathematics 1 **May/June 2020**

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages. Blank pages are indicated.

© UCLES 2020

1 The cubic equation $7x^3 + 3x^2 + 5x + 1 = 0$ has roots α, β, γ.

(a) Find a cubic equation whose roots are α^{-1}, β^{-1}, γ^{-1}. [3]

.....

.....

.....

.....

.....

.....

.....

(b) Find the value of $\alpha^{-2} + \beta^{-2} + \gamma^{-2}$. [2]

.....

.....

.....

.....

.....

.....

.....

(c) Find the value of $\alpha^{-3} + \beta^{-3} + \gamma^{-3}$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

2 The sequence $u_1, u_2, u_3, \ldots$ is such that $u_1 = 1$ and $u_{n+1} = 2u_n + 1$ for $n \geqslant 1$.

(a) Prove by induction that $u_n = 2^n - 1$ for all positive integers n. [5]

(b) Deduce that u_{2n} is divisible by u_n for $n \geqslant 1$. [2]

3 Let $S_n = 2^2 + 6^2 + 10^2 + \ldots + (4n-2)^2$.

(a) Use standard results from the List of Formulae (MF19) to show that $S_n = \frac{4}{3}n(4n^2 - 1)$. [4]

(b) Express $\frac{n}{S_n}$ in partial fractions and find $\sum_{n=1}^{N} \frac{n}{S_n}$ in terms of N. [4]

(c) Deduce the value of $\sum_{n=1}^{\infty} \frac{n}{S_n}$. [1]

4 The matrix **A** is given by

$$\mathbf{A} = \begin{pmatrix} k & 0 & 2 \\ 0 & -1 & -1 \\ 1 & 1 & -k \end{pmatrix},$$

where k is a real constant.

(a) Show that **A** is non-singular. [3]

.....

.....

.....

.....

.....

.....

.....

.....

The matrices **B** and **C** are given by

$$\mathbf{B} = \begin{pmatrix} 0 & -3 \\ -1 & 3 \\ 0 & 0 \end{pmatrix} \text{ and } \mathbf{C} = \begin{pmatrix} -3 & -1 & 1 \\ 1 & 1 & 2 \end{pmatrix}.$$

It is given that $\mathbf{CAB} = \begin{pmatrix} -2 & -\frac{3}{2} \\ -1 & -\frac{3}{2} \end{pmatrix}$.

(b) Find the value of k. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

(c) Find the equations of the invariant lines, through the origin, of the transformation in the x-y plane represented by **CAB**. [5]

5 The curve C has polar equation $r = a\tan\theta$, where a is a positive constant and $0 \leqslant \theta \leqslant \frac{1}{4}\pi$.

(a) Sketch C and state the greatest distance of a point on C from the pole. [2]

(b) Find the exact value of the area of the region bounded by C and the half-line $\theta = \frac{1}{4}\pi$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

(c) Show that C has Cartesian equation $y = \dfrac{x^2}{\sqrt{a^2 - x^2}}$. [3]

(d) Using your answer to part **(b)**, deduce the exact value of $\displaystyle\int_0^{\frac{1}{2}a\sqrt{2}} \frac{x^2}{\sqrt{a^2 - x^2}}\,\mathrm{d}x$. [2]

BLANK PAGE

© UCLES 2020

6 The curve C has equation $y = \dfrac{10 + x - 2x^2}{2x - 3}$.

(a) Find the equations of the asymptotes of C. [3]

(b) Show that C has no turning points. [3]

(c) Sketch C, stating the coordinates of the intersections with the axes. [3]

© UCLES 2020

(d) Sketch the curve with equation $y = \left|\frac{10+x-2x^2}{2x-3}\right|$ and find in exact form the set of values of x for which $\left|\frac{10+x-2x^2}{2x-3}\right| < 4$. [6]

7 The lines l_1 and l_2 have equations $\mathbf{r} = -5\mathbf{j} + \lambda(5\mathbf{i} + 2\mathbf{k})$ and $\mathbf{r} = 4\mathbf{i} + 2\mathbf{j} - 2\mathbf{k} + \mu(\mathbf{j} + \mathbf{k})$ respectively. The plane Π contains l_1 and is parallel to l_2.

(a) Find the equation of Π, giving your answer in the form $ax + by + cz = d$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the distance between l_2 and Π. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

The point P on l_1 and the point Q on l_2 are such that PQ is perpendicular to both l_1 and l_2.

(c) Show that P has position vector $\frac{55}{27}\mathbf{i} - 5\mathbf{j} + \frac{22}{27}\mathbf{k}$ and state a vector equation for PQ. [8]

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

© UCLES 2020