Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME

CENTRE NUMBER | | | | | |

CANDIDATE NUMBER | | | | |

6007163164

FURTHER MATHEMATICS **9231/33**

Paper 3 Further Mechanics **May/June 2021**

1 hour 30 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity (g) is needed, use $10\,\text{m}\,\text{s}^{-2}$.


INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages. Any blank pages are indicated.

© UCLES 2021

1



A uniform lamina $ABCD$ consists of two isosceles triangles ABD and BCD. The diagonals of $ABCD$ meet at the point O. The length of AO is $3a$, the length of OC is $6a$ and the length of BD is $16a$ (see diagram).

Find the distance of the centre of mass of the lamina from DB. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

© UCLES 2021

2 One end of a light elastic string of natural length 0.8 m and modulus of elasticity 36 N is attached to a fixed point O on a smooth plane. The plane is inclined at an angle α to the horizontal, where $\sin\alpha = \frac{3}{5}$. A particle P of mass 2 kg is attached to the other end of the string. The string lies along a line of greatest slope of the plane with the particle below the level of O. The particle is projected with speed $\sqrt{2}\,\text{m s}^{-1}$ directly down the plane from the position where OP is equal to the natural length of the string.

Find the maximum extension of the string during the subsequent motion. [5]

3




Particles A and B, of masses $3m$ and m respectively, are connected by a light inextensible string of length a that passes through a fixed smooth ring R. Particle B hangs in equilibrium vertically below the ring. Particle A moves in horizontal circles on a smooth horizontal surface with speed $\frac{2}{5}\sqrt{ga}$. The angle between AR and BR is θ (see diagram). The normal reaction between A and the surface is $\frac{12}{5}mg$.

(a) Find $\cos\theta$. [3]

(b) Find, in terms of a, the distance of B below the ring. [3]

4



A particle of mass m is attached to one end of a light inextensible string of length a. The other end of the string is attached to a fixed point O. The particle is initially held with the string taut at the point A, where OA makes an angle θ with the downward vertical through O. The particle is then projected with speed u perpendicular to OA and begins to move upwards in part of a vertical circle. The string goes slack when the particle is at the point B where angle AOB is a right angle. The speed of the particle when it is at B is $\frac{1}{2}u$ (see diagram).

Find the tension in the string at A, giving your answer in terms of m and g. [8]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

© UCLES 2021

5 A particle P of mass m kg is projected vertically upwards from a point O, with speed $20\,\text{m s}^{-1}$, and moves under gravity. There is a resistive force of magnitude $2mv$ N, where $v\,\text{m s}^{-1}$ is the speed of P at time t s after projection.

(a) Find an expression for v in terms of t, while P is moving upwards. [6]

The displacement of P from O is x m at time t s.

(b) Find an expression for x in terms of t, while P is moving upwards. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find, correct to 3 significant figures, the greatest height above O reached by P. [2]

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

6



Two uniform smooth spheres A and B of equal radii have masses m and km respectively. Sphere A is moving with speed u on a smooth horizontal surface when it collides with sphere B which is at rest. Immediately before the collision, A's direction of motion makes an angle θ with the line of centres (see diagram). The coefficient of restitution between the spheres is $\frac{1}{3}$.

(a) Show that the speed of B after the collision is $\dfrac{4u\cos\theta}{3(1+k)}$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

70% of the total kinetic energy of the spheres is lost as a result of the collision.

(b) Given that $\tan\theta = \frac{1}{3}$, find the value of k. [6]

7 A particle P is projected with speed u at an angle θ above the horizontal from a point O on a horizontal plane and moves freely under gravity. The horizontal and vertical displacements of P from O at a subsequent time t are denoted by x and y respectively.

(a) Use the equation of the trajectory given in the List of formulae (MF19), together with the condition $y = 0$, to establish an expression for the range R in terms of u, θ and g. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Deduce an expression for the maximum height H, in terms of u, θ and g. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

It is given that $R = \frac{4H}{\sqrt{3}}$.

(c) Show that $\theta = 60°$. [1]

.....

.....

.....

.....

.....

.....

It is given also that $u = \sqrt{40}\,\text{m s}^{-1}$.

(d) Find, by differentiating the equation of the trajectory or otherwise, the set of values of x for which the direction of motion makes an angle of less than 45° with the horizontal. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

BLANK PAGE

© UCLES 2021

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

© UCLES 2021