Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME

CENTRE NUMBER | | | | | |

CANDIDATE NUMBER | | | | |

2464209347

FURTHER MATHEMATICS **9231/23**

Paper 2 Further Pure Mathematics 2 **May/June 2022**

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2022

1 Find the roots of the equation $z^3 = 7\sqrt{3} - 7\mathrm{i}$, giving your answers in the form $re^{i\theta}$, where $r > 0$ and $-\pi \leqslant \theta < \pi$. [5]

2 **(a)** Find the coefficient of x^2 in the Maclaurin's series for $-\ln\cos x$. [4]

(b) Find the length of the arc of the curve with equation $y = -\ln\cos x$ from the point where $x = 0$ to the point where $x = \frac{1}{4}\pi$. [4]

3 The matrix **A** is given by

$$\mathbf{A} = \begin{pmatrix} 6 & -9 & 5 \\ 5 & -8 & 5 \\ 1 & -1 & 2 \end{pmatrix}.$$

(a) Find the eigenvalues of **A**. [4]

(b) Use the characteristic equation of $\mathbf{A}$ to show that $\mathbf{A}^{-1} = p\mathbf{A}^2 + q\mathbf{I}$, where p and q are constants to be determined. [3]

4 It is given that

$$x = -t + \tan^{-1} t \qquad \text{and} \qquad y = t + \sinh^{-1} t .$$

(a) Show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = -\dfrac{t^2 + 1 + \sqrt{t^2 + 1}}{t^2}$. [4]


(b) Find the value of $\frac{d^2y}{dx^2}$ when $t = \frac{3}{4}$. [5]

5 Find the solution of the differential equation

$$x(x+7)\frac{\mathrm{d}y}{\mathrm{d}x}+7y=x$$

for which $y=7$ when $x=1$. Give your answer in the form $y=\mathrm{f}(x)$. [9]

6



The diagram shows the curve with equation $y = \ln(1+x)$ for $0 \leqslant x \leqslant 1$, together with a set of n rectangles each of width $\frac{1}{n}$.

(a) By considering the sum of the areas of these rectangles, show that $\int_0^1 \ln(1+x)\,\mathrm{d}x < U_n$, where

$$U_n = \frac{1}{n}\ln\frac{(2n)!}{n!} - \ln n.$$ [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Use a similar method to find, in terms of n, a lower bound L_n for $\int_0^1 \ln(1+x)\,\mathrm{d}x$. [4]

(c) By simplifying $U_n - L_n$, show that $\lim_{n\to\infty}(U_n - L_n) = 0$. [2]

7 The variables x and y are related by the differential equation

$$4\frac{\mathrm{d}^2y}{\mathrm{d}x^2} - y = 3.$$

It is given that, when $x = 0$, $y = -3$ and $\frac{\mathrm{d}y}{\mathrm{d}x} = 2$.

(a) Find y in terms of x. [8]

(b) Deduce the exact value of x for which $y = 0$. Give your answer in logarithmic form. [3]

8 **(a)** Find $\int \sin\theta \cos^n\theta \, \mathrm{d}\theta$, where $n \neq -1$. [2]

.....

.....

.....

Let $I_{m,n} = \int_0^{\frac{1}{2}\pi} \sin^m\theta \cos^n\theta \, \mathrm{d}\theta$.

(b) Show that, for $m \geqslant 2$ and $n \geqslant 0$,

$$I_{m,n} = \frac{m-1}{m+n} I_{m-2,n}.$$ [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

(c) By considering the binomial expansion of $\left(z+\frac{1}{z}\right)^5$, where $z=\cos\theta+\mathrm{i}\sin\theta$, use de Moivre's theorem to show that

$$\cos^5\theta = a\cos 5\theta + b\cos 3\theta + c\cos\theta,$$

where a, b and c are constants to be determined. [5]

(d) Using the results given in parts **(b)** and **(c)**, find the exact value of $I_{2,5}$. [4]

Additional page

If you use the following page to complete the answer to any question, the question number must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

BLANK PAGE

© UCLES 2022

BLANK PAGE

© UCLES 2022

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2022