Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME

CENTRE NUMBER | | | | | |

CANDIDATE NUMBER | | | | |

0834557711

FURTHER MATHEMATICS **9231/13**

Paper 1 Further Pure Mathematics 1 **May/June 2023**

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages. Any blank pages are indicated.

© UCLES 2023

BLANK PAGE

© UCLES 2023

1 Prove by mathematical induction that, for all positive integers n, $5^{3n} + 32^n - 33$ is divisible by 31. [6]

2 **(a)** Use standard results from the list of formulae (MF19) to show that

$$\sum_{r=1}^{n}(6r^2+6r-5) = an^3+bn^2+cn,$$

where a, b and c are integers to be determined. [2]

(b) Use the method of differences to find $\sum_{r=1}^{n}\frac{6r^2+6r-5}{r^2+r}$ in terms of n. [4]

(c) Find also $\sum_{r=n+1}^{2n} \frac{6r^2+6r-5}{r^2+r}$ in terms of n. [2]

3 The equation $x^4 - x^2 + 2x + 5 = 0$ has roots $\alpha, \beta, \gamma, \delta$.

(a) Find a quartic equation whose roots are $\alpha^2, \beta^2, \gamma^2, \delta^2$ and state the value of $\alpha^2 + \beta^2 + \gamma^2 + \delta^2$. [4]

(b) Find the value of $\frac{1}{\alpha^2}+\frac{1}{\beta^2}+\frac{1}{\gamma^2}+\frac{1}{\delta^2}$. [3]

(c) Find the value of $\alpha^4+\beta^4+\gamma^4+\delta^4$. [2]

4 The matrix $\mathbf{M}$ is given by $\mathbf{M} = \begin{pmatrix} \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{pmatrix} \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}$, where $0 < \theta < \pi$ and k is a non-zero constant. The matrix $\mathbf{M}$ represents a sequence of two geometrical transformations, one of which is a shear.

(a) Describe fully the other transformation and state the order in which the transformations are applied. [3]

.....

.....

.....

.....

.....

.....

(b) Write $\mathbf{M}^{-1}$ as the product of two matrices, neither of which is $\mathbf{I}$. [2]

.....

.....

.....

.....

.....

.....

(c) Find, in terms of k, the value of $\tan\theta$ for which $\mathbf{M} - \mathbf{I}$ is singular. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

(d) Given that $k = 2\sqrt{3}$ and $\theta = \frac{1}{3}\pi$, show that the invariant points of the transformation represented by **M** lie on the line $3y + \sqrt{3}x = 0$. [4]

5 **(a)** Show that the curve with Cartesian equation

$$x^2 - y^2 = a,$$

where a is a positive constant, has polar equation $r^2 = a\sec 2\theta$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

The curve C has polar equation $r^2 = a\sec 2\theta$, where a is a positive constant, for $0 \leqslant \theta < \frac{1}{4}\pi$.

(b) Sketch C and state the minimum distance of C from the pole. [3]

.....

(c) Find, in terms of a, the exact value of the area of the region enclosed by C, the initial line, and the half-line $\theta = \frac{1}{12}\pi$. [You may use any result from the list of formulae (MF19) without proof.] [4]

6 The points A, B, C have position vectors

$$\mathbf{i}+\mathbf{j}, \quad -\mathbf{i}+2\mathbf{j}+4\mathbf{k}, \quad -2\mathbf{i}+\mathbf{j}+3\mathbf{k},$$

respectively, relative to the origin O.

(a) Find the equation of the plane ABC, giving your answer in the form $ax+by+cz=d$. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the perpendicular distance from O to the plane ABC. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

(c) Find a vector equation of the common perpendicular to the lines OC and AB. [8]

7 The curve C has equation $y = \dfrac{x^2 + 2x + 1}{x - 3}$.

(a) Find the equations of the asymptotes of C. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the coordinates of the turning points on C. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

(c) Sketch C. [3]

(d) Sketch the curves with equations $y = \left| \frac{x^2 + 2x + 1}{x - 3} \right|$ and $y^2 = \frac{x^2 + 2x + 1}{x - 3}$ on a single diagram, clearly identifying each curve. [4]

© UCLES 2023

Additional page

If you use the following page to complete the answer to any question, the question number must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2023