Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME

CENTRE NUMBER

CANDIDATE NUMBER

7729284446

FURTHER MATHEMATICS **9231/22**

Paper 2 Further Pure Mathematics 2 **October/November 2020**

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages. Blank pages are indicated.

DC (RW/SG) 187470/2
© UCLES 2020

[Turn over

1 Find the Maclaurin's series for $\tan\left(x+\frac{1}{4}\pi\right)$ up to and including the term in x^2. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

2 A curve has equation $y = \cosh x$, for $0 \leqslant x \leqslant \frac{1}{2}$.

Find, in terms of π and e, the area of the surface generated when the curve is rotated through 2π radians about the x-axis. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3 Find all the roots of the equation $(w+1)^6 = 1$, giving your answers in the form $x+\mathrm{i}y$ where x and y are real and exact. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 Find the solution of the differential equation

$$x\frac{\mathrm{d}y}{\mathrm{d}x} + 2y = \mathrm{e}^x$$

for which $y = 3$ when $x = 1$. Give your answer in the form $y = \mathrm{f}(x)$. [8]

5 The curve C has equation

$$y^2 + (xy+1)^2 = 5.$$

(a) Show that, at the point $(1,1)$ on C, $\frac{\mathrm{d}y}{\mathrm{d}x} = -\frac{2}{3}$. [3]

(b) Find the value of $\frac{\mathrm{d}^2y}{\mathrm{d}x^2}$ at the point $(1,1)$. [5]

6 Find the particular solution of the differential equation

$$\frac{\mathrm{d}^2x}{\mathrm{d}t^2} + 8\frac{\mathrm{d}x}{\mathrm{d}t} + 15x = 102\cos 3t,$$


given that, when $t = 0$, $x = 1$ and $\frac{\mathrm{d}x}{\mathrm{d}t} = 0$. [11]

7 **(a)** Show that $\sum_{r=1}^{n} z^{2r} = \frac{z^{2n+1} - z}{z - z^{-1}}$, for $z \neq 0, 1, -1$. [2]

(b) By letting $z = \cos\theta + \mathrm{i}\sin\theta$, show that, if $\sin\theta \neq 0$,

$$1 + 2\sum_{r=1}^{n} \cos(2r\theta) = \frac{\sin(2n+1)\theta}{\sin\theta}. \qquad [5]$$

8



The diagram shows the curve $y = \dfrac{1}{\sqrt{x^2+x+1}}$ for $x \geqslant 0$, together with a set of n rectangles of unit width. By considering the sum of the areas of these rectangles, show that

$$\sum_{r=1}^{n} \frac{1}{\sqrt{r^2+r+1}} < \ln\left(\tfrac{1}{3} + \tfrac{2}{3}n + \tfrac{2}{3}\sqrt{n^2+n+1}\right). \qquad [10]$$

9 It is given that a is a positive constant.

(a) Show that the system of equations

$$\begin{aligned} ax + (2a+5)y + (a+1)z &= 1, \\ -4y \qquad\quad &= 2, \\ 3y - z &= 3, \end{aligned}$$

has a unique solution and interpret this situation geometrically. [3]

The matrix **A** is given by

$$\mathbf{A} = \begin{pmatrix} a & 2a+5 & a+1 \\ 0 & -4 & 0 \\ 0 & 3 & -1 \end{pmatrix}.$$

(b) Show that the eigenvalues of **A** are a, -1 and -4. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find a matrix **P** such that

$$\mathbf{A} = \mathbf{P}\begin{pmatrix} a & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -4 \end{pmatrix}\mathbf{P}^{-1}.$$ [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

(d) Use the characteristic equation of $\mathbf{A}$ to find $\mathbf{A}^{-1}$. [6]

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

© UCLES 2020