Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

6087072568

FURTHER MATHEMATICS **9231/13**

Paper 1 Further Pure Mathematics 1 **October/November 2022**

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages. Any blank pages are indicated.

© UCLES 2022

1 The cubic equation $x^3 + bx^2 + d = 0$ has roots α, β, γ, where $\alpha = \beta$ and $d \neq 0$.

(a) Show that $4b^3 + 27d = 0$. [5]

(b) Given that $2\alpha^2 + \gamma^2 = 3b$, find the values of b and d. [3]

.....

.....

.....

.....

.....

2 Prove by mathematical induction that, for all positive integers n, $7^{2n} + 97^n - 50$ is divisible by 48. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3 **(a)** By considering $(2r+1)^3-(2r-1)^3$, use the method of differences to prove that

$$\sum_{r=1}^{n} r^2 = \tfrac{1}{6}n(n+1)(2n+1).$$ [5]

Let $S_n = 1^2 + 3\times 2^2 + 3^2 + 3\times 4^2 + 5^2 + 3\times 6^2 + \ldots + \left(2 + (-1)^n\right)n^2$.

(b) Show that $S_{2n} = \frac{1}{3}n(2n+1)(an+b)$, where a and b are integers to be determined. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) State the value of $\lim_{n\to\infty} \frac{S_{2n}}{n^3}$. [1]

.....

.....

.....

.....

.....

.....

.....

4 The plane Π contains the lines $\mathbf{r} = 3\mathbf{i} - 2\mathbf{j} + \mathbf{k} + \lambda(-\mathbf{i} + 2\mathbf{j} + \mathbf{k})$ and $\mathbf{r} = 4\mathbf{i} + 4\mathbf{j} + 2\mathbf{k} + \mu(3\mathbf{i} + 2\mathbf{j} - \mathbf{k})$.

(a) Find a Cartesian equation of Π, giving your answer in the form $ax + by + cz = d$. [4]

The line l passes through the point P with position vector $2\mathbf{i}+3\mathbf{j}+\mathbf{k}$ and is parallel to the vector $\mathbf{j}+\mathbf{k}$.

(b) Find the acute angle between l and Π. [3]

(c) Find the position vector of the foot of the perpendicular from P to Π. [4]

5 The matrix **M** is given by $\mathbf{M} = \begin{pmatrix} \frac{1}{2}\sqrt{2} & -\frac{1}{2}\sqrt{2} \\ \frac{1}{2}\sqrt{2} & \frac{1}{2}\sqrt{2} \end{pmatrix}\begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}$, where k is a constant.

(a) The matrix **M** represents a sequence of two geometrical transformations.

State the type of each transformation, and make clear the order in which they are applied. [2]

.....

.....

.....

.....

.....

.....

.....

.....

(b) The triangle ABC in the x–y plane is transformed by **M** onto triangle DEF.

Find, in terms of k, the single matrix which transforms triangle DEF onto triangle ABC. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

(c) Find the set of values of k for which the transformation represented by **M** has no invariant lines through the origin. [7]

6 **(a)** Show that the curve with Cartesian equation

$$\left(x^2+y^2\right)^2 = 36\left(x^2-y^2\right)$$

has polar equation $r^2 = 36\cos 2\theta$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

The curve C has polar equation $r^2 = 36\cos 2\theta$, for $-\frac{1}{4}\pi \leqslant \theta \leqslant \frac{1}{4}\pi$.

(b) Sketch C and state the maximum distance of a point on C from the pole. [3]

.....

.....

© UCLES 2022

(c) Find the area of the region enclosed by C. [2]

(d) Find the maximum distance of a point on C from the initial line, giving the answer in exact form. [6]

7 The curve C has equation $y = \dfrac{5x^2}{5x-2}$.

(a) Find the equations of the asymptotes of C. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the coordinates of the stationary points on C. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Sketch C. [3]

(d) Sketch the curve with equation $y = \left|\frac{5x^2}{5x-2}\right|$ and find in exact form the set of values of x for which $\left|\frac{5x^2}{5x-2}\right| < 2$. [6]

© UCLES 2022

Additional page

If you use the following page to complete the answer to any question, the question number must be clearly shown.

© UCLES 2022

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2022