UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education
Advanced Subsidiary Level and Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

3576889954

CHEMISTRY **9701/22**

Paper 2 Structured Questions AS Core **October/November 2013**

1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials: Data Booklet

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use a soft pencil for any diagrams, graphs or rough working.
Do not use staples, paper clips, highlighters, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.
Electronic calculators may be used.
You may lose marks if you do not show your working or if you do not use appropriate units.
A Data Booklet is provided.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.

For Examiner's Use	
1	
2	
3	
4	
5	
Total	

This document consists of **9** printed pages and **3** blank pages.

© UCLES 2013

UNIVERSITY of CAMBRIDGE International Examinations

Answer **all** the questions in the spaces provided.

For Examiner's Use

1 Valence Shell Electron Pair Repulsion theory (VSEPR) is a model of electron-pair repulsion (including lone pairs) that can be used to deduce the shapes of, and bond angles in, simple molecules.

(a) Complete the table below by using simple hydrogen-containing compounds. One example has been included.

number of bond pairs	number of lone pairs	shape of molecule	formula of a molecule with this shape
3	0	trigonal planar	BH_3
4	0		
3	1		
2	2		

[3]

(b) Tellurium, Te, proton number 52, is used in photovoltaic cells.

When fluorine gas is passed over tellurium at 150 °C, the colourless gas TeF_6 is formed.

(i) Draw a 'dot-and-cross' diagram of the TeF_6 molecule, showing outer electrons only.

(ii) What will be the shape of the TeF_6 molecule?

.....

(iii) What is the F–Te–F bond angle in TeF_6?

.....


[3]

[Total: 6]


© UCLES 2013

For Examiner's Use

2 The molecular formula C_3H_6 represents the compounds propene and cyclopropane.



propene



cyclopropane

(a) What is the H–C–H bond angle at the terminal $=CH_2$ group in propene?

..... [1]

(b) Under suitable conditions, propene and cyclopropane each react with chlorine.

(i) With propene, 1,2-dichloropropane, $CH_3CHCl CH_2Cl$ is formed.

State fully what type of reaction this is.

..... [1]

(ii) When cyclopropane reacts with chlorine, three different compounds with the molecular formula $C_3H_4Cl_2$ can be formed.

Draw displayed structures of **each** of these three compounds.

[3]

[Total: 5]

For Examiner's Use

3 Chlorine gas is manufactured by the electrolysis of brine using a diaphragm cell.

(a) (i) Write half-equations, including state symbols, for the reactions occurring at **each** of the electrodes of a diaphragm cell.

anode .....

cathode .....

(ii) In the diaphragm cell, the anode is made of titanium and the cathode is made of steel.

Suggest why steel is never used for the anode.

.....

..... [3]

(b) Chlorine is very reactive and will form compounds by direct combination with many elements.

Describe what you would see when chlorine is passed over separate heated samples of sodium and phosphorus.
In **each** case write an equation for the reaction.

sodium

.....

.....

.....

phosphorus

.....

.....

..... [4]

© UCLES 2013

For Examiner's Use

(c) Chlorine reacts with aqueous sodium hydroxide in two different ways, depending on the conditions used. In each case, water, sodium chloride and one other chlorine-containing compound are formed.

For **each** condition below, give the formula of the **other** chlorine-containing compound and state the oxidation number of chlorine in it.

condition	formula of **other** chlorine-containing compound	oxidation number of chlorine in this compound
cold dilute NaOH(aq)		
hot concentrated NaOH(aq)		

[4]

(d) Magnesium chloride, $MgCl_2$, and silicon tetrachloride, $SiCl_4$, each dissolve in or react with water.

Suggest the approximate pH of the solution formed in **each** case.

$MgCl_2$ ..... $SiCl_4$ .....

Explain, with the aid of an equation, the difference between the two values.

.....

.....

.....

..... [5]

[Total: 16]

© UCLES 2013

[Turn over

For Examiner's Use

4 Compound **R** is a weak diprotic (dibasic) acid which is very soluble in water.

(a) A solution of **R** was prepared which contained 1.25 g of **R** in 250 cm^3 of solution. When 25.0 cm^3 of this solution was titrated with 0.100 mol dm^{-3} NaOH, 21.6 cm^3 of the alkali were needed for complete reaction.

(i) Using the formula H_2X to represent **R**, construct a balanced equation for the reaction between H_2X and NaOH.

.....

(ii) Use the data above to calculate the amount, in moles, of OH^- ions used in the titration.

(iii) Use your answers to **(i)** and **(ii)** to calculate the amount, in moles, of **R** present in 25.0 cm^3 of solution.

(iv) Calculate the amount, in moles, of **R** present in 250 cm^3 of solution.

(v) Calculate M_r of **R**.

[5]

(b) Three possible structures for **R** are shown below.

S	**T**	**U**
$HO_2CCH{=}CHCO_2H$	$HO_2CCH(OH)CH_2CO_2H$	$HO_2CCH(OH)CH(OH)CO_2H$

(i) Calculate the M_r of each of these acids.

M_r of **S** = ... M_r of **T** = ... M_r of **U** = ...

(ii) Deduce which of the structures, **S**, **T** or **U**, correctly represents the structure of the acid, **R**.

R is represented by .....

[2]

© UCLES 2013

It is possible to convert **S**, **T**, or **U** into one another.

(c) State the reagent(s) and essential conditions that would be used for the following conversions.

S into **T**

.....

S into **U**

.....

T into **S**

..... [5]

(d) Give the structural formula of the organic product formed in **each** of the following reactions.

T reacting with an excess of Na

U reacting with an excess of Na_2CO_3

[2]

(e) The acid **S** shows stereoisomerism. Draw structures to show this isomerism.
Label each isomer.

[2]

(f) When one of the isomers of **S** is heated at 110 °C in the absence of air, a cyclic compound **V**, with molecular formula $C_4H_2O_3$, is formed.
The other isomer of **S** does not react at this temperature.

Suggest the displayed formula of **V**.

[2]

[Total: 18]

© UCLES 2013

For Examiner's Use

5 Propane, C_3H_8, and butane, C_4H_{10}, are components of Liquefied Petroleum Gas (LPG) which is widely used as a fuel for domestic cooking and heating.

(a) (i) To which class of compounds do these two hydrocarbons belong?

.....

(ii) Write a balanced equation for the complete combustion of butane.

.....
[2]

(b) When propane or butane is used in cooking, the saucepan may become covered by a solid black deposit.

(i) What is the chemical name for this black solid?

.....

(ii) Write a balanced equation for its formation from butane.

.....
[2]

(c) Propane and butane have different values of standard enthalpy change of combustion.

Define the term *standard enthalpy change of combustion*.

.....

.....

..... [2]

(d) A 125 cm^3 sample of propane gas, measured at 20 °C and 101 kPa, was completely burnt in air.
The heat produced raised the temperature of 200 g of water by 13.8 °C.
Assume no heat losses occurred during this experiment.

(i) Use the equation $pV = nRT$ to calculate the mass of propane used.

© UCLES 2013

(ii) Use relevant data from the *Data Booklet* to calculate the amount of heat released in this experiment.

(iii) Use the data above and your answers to **(i)** and **(ii)** to calculate the energy produced by the burning of 1 mol of propane.

[5]

(e) The boiling points of methane, ethane, propane, and butane are given below.

compound	CH_4	CH_3CH_3	$CH_3CH_2CH_3$	$CH_3(CH_2)_2CH_3$
boiling point / K	112	185	231	273

(i) Suggest an explanation for the increase in boiling points from methane to butane.

.....

.....

.....

(ii) The isomer of butane, 2-methylpropane, $(CH_3)_3CH$, has a boiling point of 261 K. Suggest an explanation for the difference between this value and that for butane in the table above.

.....

.....

.....

[4]

[Total: 15]

© UCLES 2013

BLANK PAGE

© UCLES 2013

BLANK PAGE

© UCLES 2013

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.