CAMBRIDGE
International Education

Cambridge International AS & A Level

CHEMISTRY **9701/22**

Paper 2 AS Structured Questions **October/November 2024**

MARK SCHEME

Maximum Mark: 60

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2024 series for most Cambridge IGCSE, Cambridge International A and AS Level components, and some Cambridge O Level components.

© Cambridge University Press & Assessment 2024

PUBLISHED

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptions for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

© Cambridge University Press & Assessment 2024

GENERIC MARKING PRINCIPLE 5: Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).
GENERIC MARKING PRINCIPLE 6: Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Science-Specific Marking Principles

1	Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.
2	The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.
3	Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).
4	The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.
5	‘List rule’ guidance For questions that require ***n*** responses (e.g. State **two** reasons …): • The response should be read as continuous prose, even when numbered answer spaces are provided. • Any response marked *ignore* in the mark scheme should not count towards ***n***. • Incorrect responses should not be awarded credit but will still count towards ***n***. • Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should **not** be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response. • Non-contradictory responses after the first ***n*** responses may be ignored even if they include incorrect science.

© Cambridge University Press & Assessment 2024

6 Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form (e.g. $a \times 10^n$) in which the convention of restricting the value of the coefficient (a) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

7 Guidance for chemical equations

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.

© Cambridge University Press & Assessment 2024

<table>
<tr><th>Question</th><th>Answer</th><th>Marks</th></tr>
<tr><td>1(a)(i)</td><td>lowest energy state owtte

examples
• (with) lowest energy
• (with) no external energy
• (with) no added energy
• (where) electrons are not promoted to a higher energy level
• most energetically stable</td><td>1</td></tr>
<tr><td>1(a)(ii)</td><td>[Ar] 
The 3 × d electrons may go in any of the d orbitals</td><td>1</td></tr>
<tr><td>1(a)(iii)</td><td>12</td><td>1</td></tr>
<tr><td>1(b)(i)</td><td><table><tr><th>isotope</th><th>RIM</th><th># protons</th><th># neutrons</th></tr><tr><td>$^{93}_{41}$Nb</td><td>92.91</td><td>● 41</td><td>● 52</td></tr><tr><td>$^{181}_{73}$Ta</td><td>180.95</td><td>● 73</td><td>● 108</td></tr></table></td><td>2</td></tr>
</table>

© Cambridge University Press & Assessment 2024

Question	Answer	Marks
1(b)(ii)	**M1** mass of an (atom of an) isotope	**1**
	M2 relative/compared to (the mass of) the unified atomic mass unit **OR** on a scale in which a carbon-12 atom / isotope has a mass of exactly 12 units OR divided by / compared to $\frac{1}{12}$ of the mass of a carbon-12 atom / isotope Alternative route using mass of 1 mol throughout **M1** mass of one mol of an (atom of an) isotope **M2** relative/compared to $\frac{1}{12}$ of the mass of 1 mol of C-12 atom / isotope **OR** when the mass of one mol C-12 atom / isotope is exactly 12(.000) g	**1**
1(b)(iii)	$92.91 \times 0.909 + 180.95 \times 0.091$	**1**
	= 100.92	**1**

Question	Answer	Marks
2(a)(i)	M1 $4Na + O_2 \rightarrow 2Na_2O$	**1**
	M2 $S + O_2 \rightarrow SO_2$	**1**
2(a)(ii)	Al^{3+} 	**1**
2(a)(iii)	it increases	**1**
	as number of valence electrons (which can be used in bonding / lost / shared / donated) increases	**1**

© Cambridge University Press & Assessment 2024

<table>
<tr><th>Question</th><th>Answer</th><th>Marks</th></tr>
<tr><td>2(b)(i)</td><td>
<table>
<tr><td>Period 3 oxide</td><td>product of rxn with water</td><td>pH of solution formed</td></tr>
<tr><td>● MgO</td><td>$Mg(OH)_2$</td><td>● $8 \leqslant pH \leqslant 12$</td></tr>
<tr><td>P_4O_{10}</td><td>● H_3PO_4</td><td>● $1 \leqslant pH \leqslant 4$</td></tr>
</table>
●✓●✓
</td><td>2</td></tr>
<tr><td>2(b)(ii)</td><td>$CH_3CN + 2H_2O + (1)H^+ \rightarrow CH_3COOH + NH_4^+$</td><td>1</td></tr>
<tr><td>2(b)(iii)</td><td></td><td>2</td></tr>
<tr><td rowspan="2">2(b)(iv)</td><td>alkyl groups are electron donating / have positive inductive effect.</td><td>1</td></tr>
<tr><td>strengthens the O—H bond / makes H^+ less likely to be donated</td><td>1</td></tr>
<tr><td rowspan="2">2(c)(i)</td><td>(molecules) H_2S to H_2Te / they have greater number of electrons</td><td>1</td></tr>
<tr><td>stronger instantaneous dipole–induced dipole / London dispersion forces
OR
more energy required to overcome the instantaneous dipole–induced dipole / London dispersion forces</td><td>1</td></tr>
<tr><td>2(c)(ii)</td><td>(only) H_2O has hydrogen bonding AND hydrogen bonding (much) stronger than the other intermolecular forces
OR
H_2O has hydrogen bonding AND because of higher electronegativity of O compared to S
OR
stronger van der waals’ forces because (only) H_2O has hydrogen bonding
OR
hydrogen bond AND increased strength of permanent dipoles in H_2O outweighs the increase in strength of id-id in the others</td><td>1</td></tr>
</table>




© Cambridge University Press & Assessment 2024

Question	Answer	Marks
3(a)(i)	(natural =)lightning	**1**
	(man-made =)internal combustion engines	**1**
3(a)(ii)	$2NO_2 + H_2O \rightarrow HNO_2 + HNO_3$ **OR** $4NO_2 + O_2 + 2H_2O \rightarrow 4HNO_3$	**1**
3(a)(iii)	It / NO_2 reacts with (unburned) hydrocarbons / VOCs **ALLOW** reaction of unburned hydrocarbons / VOCs in presence of NO_2	**1**
3(a)(iv)	$2HNO_3 + CaO \rightarrow Ca(NO_3)_2 + H_2O$	**1**
3(a)(v)	brown fumes given off	**1**
3(b)(i)	(+)5 / V	**1**
3(b)(ii)	aluminium / Al	**1**
3(b)(iii)	(NH_3 is an) H^+ acceptor	**1**
3(b)(iv)	tetrahedral	**1**
3(c)(i)	$P^{2+}(g) \rightarrow P^{3+}(g) + e^{(-)}$	**1**
3(c)(ii)	general increase	**1**
	increase in IE between IE5 and IE6 is noticeably the largest increase between IEs shown	**1**


© Cambridge University Press & Assessment 2024

<table>
<tr><th>Question</th><th>Answer</th><th>Marks</th></tr>
<tr><td>3(d)</td><td>
<table>
<tr><td>element</td><td>nitrogen</td><td>phosphorus</td></tr>
<tr><td>state and appearance</td><td>colourless gas</td><td>white solid</td></tr>
<tr><td>electrical conductivity</td><td>● poor</td><td>poor</td></tr>
<tr><td>type of bonding</td><td>● covalent</td><td>● covalent</td></tr>
<tr><td>type of structure</td><td>simple</td><td>● simple</td></tr>
</table>
</td><td>2</td></tr>
<tr><td>3(e)</td><td>simple molecular (lattice structure)</td><td>1</td></tr>
<tr><td rowspan="2">3(f)(i)</td><td>M1 one sigma / σ bond AND head-on (overlap of) p / sp (orbitals)</td><td>1</td></tr>
<tr><td>M2 two pi / π bond(s) AND side-on (overlap of / involving) p (orbitals)</td><td>1</td></tr>
<tr><td>3(f)(ii)</td><td>P≡P is much weaker so P_2 is more reactive (than N_2)</td><td>1</td></tr>
</table>

© Cambridge University Press & Assessment 2024

Question	Answer	Marks
4(a)	 **pt 1** dipole on Br_2 **pt 2** curly arrow from C=C bond to a $Br^{\delta+}$ **pt 3** curly arrow from Br–Br bond to the other Br **pt 4** correct intermediate **pt 5** curly arrow from lone pair on Br^- to C^+ **5•** = 3 marks. **4•** = 2 marks. **3•** = 2 marks. **2•** = 1 marks. **1•** = 0 marks.	**3**
4(b)	= –37.8 (kJ mol^{-1})	**1**
4(c)(i)		**1**
	bromoethene	**1**
4(c)(ii)	 OR	**1**
4(c)(iii)	C_2H_2	**1**
4(d)(i)	structural / positional	**1**
4(d)(ii)	aldehyde	**1**

© Cambridge University Press & Assessment 2024

<table>
<tr><th>Question</th><th>Answer</th><th>Marks</th></tr>
<tr><td>4(d)(iii)</td><td>
<table>
<tr><td>reagent</td><td>observation with F</td></tr>
<tr><td>2,4-DNPH</td><td>red / orange / yellow ppt</td></tr>
<tr><td>Tollens' reagent</td><td>silver mirror OR grey ppt OR black ppt</td></tr>
<tr><td>alkaline I_2(aq)</td><td>yellow ppt</td></tr>
</table>
</td><td>3</td></tr>
<tr><td>4(e)(i)</td><td>M1 identify H = 

M2 (broad) absorption within the range 3600–2500 cm^{-1} so O-H (bond)
O-H (bond) is equivalent to OH <u>bond</u>

M3 pt 1 and pt 2 •✓ OR pt 1 and pt 3 •✓

•pt 1 absorption within the range 1670–1750 cm^{-1} so C=O (bond)

•pt 2 absorption within the range 1040–1300 cm^{-1} so C-O (bond)

•pt 3 (M^+ at m / e = 60 so it has) molecular mass / Mr = 60</td><td></td></tr>
<tr><td>4(e)(ii)</td><td>oxidising agent</td><td>1</td></tr>
</table>

© Cambridge University Press & Assessment 2024