* 0000800000001 * DFD

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME

CENTRE NUMBER

CANDIDATE NUMBER

4215058992

CHEMISTRY **9701/34**

Paper 3 Advanced Practical Skills 2 **October/November 2025**

2 hours

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [ ].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.
- Notes for use in qualitative analysis are provided in the question paper.

Session
Laboratory

For Examiner's Use	
1	
2	
3	
Total	

This document has **12** pages.

© UCLES 2025

[Turn over

Quantitative analysis

Read through the whole method before starting any practical work. Where appropriate, prepare a table for your results in the space provided.

Show the precision of the apparatus you used in the data you record.

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

1 Sodium carbonate can be manufactured using a two-step process. The first step involves making sodium hydrogencarbonate, $NaHCO_3$, which is then converted into sodium carbonate, Na_2CO_3, in the second step of the process by the following reaction.

$$2NaHCO_3(s) \rightarrow Na_2CO_3(s) + H_2O(l) + CO_2(g)$$

In this experiment you will determine the enthalpy change, ΔH_r, for this reaction. You will do this by calculating the enthalpy changes when separate samples of sodium hydrogencarbonate and sodium carbonate are added to excess hydrochloric acid, HC*l*(aq). You will then combine these values using Hess's law.

$$NaHCO_3(s) + HCl(aq) \rightarrow NaCl(aq) + H_2O(l) + CO_2(g) \qquad \Delta H_1$$

$$Na_2CO_3(s) + 2HCl(aq) \rightarrow 2NaCl(aq) + H_2O(l) + CO_2(g) \qquad \Delta H_2$$

FB 1 is 2.0 mol dm^{-3} hydrochloric acid, HC*l*.
FB 2 is sodium hydrogencarbonate, $NaHCO_3$.
FB 3 is sodium carbonate, Na_2CO_3.

(a) Method

Experiment 1

- Support a cup in the 250 cm^3 beaker.
- Use the 25 cm^3 measuring cylinder to transfer 20.0 cm^3 of **FB 1** into the cup.
- Weigh the container with **FB 2**. Record the mass.
- Measure the temperature of the acid in the cup. Record this temperature.
- Carefully add all of the **FB 2**, in small portions to avoid acid spray. Stir to dissolve. Record the lowest temperature.
- Reweigh the container with any residual **FB 2**. Record the mass.
- Calculate and record the mass of **FB 2** used.
- Calculate and record the decrease in temperature.

Experiment 2

- Support the second cup in the 250 cm^3 beaker.
- Use the 25 cm^3 measuring cylinder to transfer 20.0 cm^3 of **FB 1** into the second cup.
- Weigh the container with **FB 3**. Record the mass.
- Measure the temperature of the acid in the cup. Record this temperature.
- Carefully add all of the **FB 3**, in small portions to avoid acid spray. Stir to dissolve. Record the highest temperature.
- Reweigh the container with any residual **FB 3**. Record the mass.
- Calculate and record the mass of **FB 3** used.
- Calculate and record the increase in temperature.

Keep FB 1 for use in Question 3.

DO NOT WRITE IN THIS MARGIN

Results

I	
II	
III	
IV	

[4]

(b) Calculations

(i) Calculate the amount, in mol, of **FB 2** that reacts with **FB 1** and the amount, in mol, of **FB 3** that reacts with **FB 1**.

Experiment 1	**Experiment 2**
amount of **FB 2** = ... mol	amount of **FB 3** = ... mol

[2]

(ii) Calculate the energy changes, in J, for each reaction.

Experiment 1	**Experiment 2**
energy change = ... J	energy change = ... J

[1]

(iii) Calculate the enthalpy change, ΔH, in $kJ\,mol^{-1}$ for each reaction.

Experiment 1	**Experiment 2**
ΔH_1 = ... $kJ\,mol^{-1}$ *sign* *value*	ΔH_2 = ... $kJ\,mol^{-1}$ *sign* *value*

[2]

DO NOT WRITE IN THIS MARGIN

(iv) Construct an enthalpy cycle and use Hess's law to determine the enthalpy change, ΔH_r, for the reaction shown.
Show your working.

$$2NaHCO_3(s) \rightarrow Na_2CO_3(s) + H_2O(l) + CO_2(g)$$

(If you were unable to calculate values in **(b)(iii)** then assume that ΔH_1 is +27.3 kJ mol^{-1} and that ΔH_2 is –24.9 kJ mol^{-1}. These may **not** be the correct values.)

ΔH_r = ... kJ mol^{-1}
sign *value*

[2]

[Total: 11]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

2 In this experiment you will determine the percentage by mass of a sodium halide impurity present in a sample of sodium hydrogencarbonate by titration.

$$NaHCO_3(s) + HCl(aq) \rightarrow NaCl(aq) + H_2O(l) + CO_2(g)$$

FB 4 is an aqueous solution made by dissolving 17.20 g of the impure sodium hydrogencarbonate in each dm^3 of solution.
FB 5 is 0.200 mol dm^{-3} hydrochloric acid, HC*l*.
FB 6 is methyl orange indicator.

(a) Method

- Fill the burette with **FB 5**.
- Pipette 25.0 cm^3 of **FB 4** into a conical flask.
- Add several drops of **FB 6** to the conical flask.
- Perform a rough titration and record your burette readings in the space below.

The rough titre is ..... cm^3.

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make sure any recorded results show the precision of your practical work.
- Record all your burette readings and the volume of **FB 5** added in each accurate titration.

Keep FB 4 for use in Question 3.

Results

I	
II	
III	
IV	
V	
VI	
VII	

[7]

(b) From your accurate titration results, calculate a suitable mean value to be used in your calculations.
Show clearly how you obtained this value.

25.0 cm^3 of **FB 4** required ..... cm^3 of **FB 5**. [1]

(c) Calculations

(i) Give your answers to **(c)(ii)** and **(c)(iii)** to an appropriate number of significant figures. [1]

(ii) Use your answer to **(b)** to calculate the amount, in mol, of hydrochloric acid in your mean titre.

amount of HCl = ... mol

Hence determine the amount, in mol, of sodium hydrogencarbonate present in 25.0 cm^3 of **FB 4**.

amount of $NaHCO_3$ = ... mol [1]

(iii) Calculate the mass of sodium hydrogencarbonate present in each dm^3 of solution.

mass of $NaHCO_3$ = ... g

Hence calculate the percentage by mass of the sodium halide impurity in **FB 4**.
Show your working.

percentage by mass = ... % [2]

(d) A student carries out the experiment in Question 1 using the impure sodium hydrogencarbonate dissolved to make **FB 4**.
State how this affects the value of ΔH_1 determined in **1(b)(iii)** compared to the value the student would get if pure sodium hydrogencarbonate were used.

.....

.....

State what assumption you have made about the sodium halide impurity.

.....

..... [2]

[Total: 14]

DO NOT WRITE IN THIS MARGIN

Qualitative analysis

For each test you should record all your observations in the spaces provided.

Examples of observations include:
- colour changes seen
- the formation of any precipitate and its solubility (where appropriate) in an excess of the reagent added
- the formation of any gas and its identification (where appropriate) by a suitable test.

You should record clearly at what stage in a test an observation is made.

Where no change is observed, you should write 'no change'.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.

If any solution is warmed, a boiling tube must be used. If any solid is heated a hard-glass test-tube must be used.

Rinse and reuse test-tubes and boiling tubes where possible.

No additional tests should be attempted.

3 (a) (i) To a 1 cm depth of **FB 4** in a boiling tube, slowly add a 1 cm depth of dilute nitric acid. Stir gently until the reaction is complete. Use this solution for your tests.
Select reagents to identify which sodium halide is present.
Record details of the reagents used and your observations.
Identify the sodium halide that is present as an impurity in **FB 4**.

The formula of the impurity is .....

[2]

(ii) Suggest why it is necessary to add the nitric acid to **FB 4** before carrying out your tests in **(a)(i)**.

.....

..... [1]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

(b) **FB 7** contains a metal cation which is listed in the Qualitative analysis notes.

(i) Place half the sample of **FB 7** in a hard-glass test-tube. Heat gently at first and then more strongly.
Record your observations.

.....

.....

..... [2]

(ii) Use the 25 cm^3 measuring cylinder to measure 20 cm^3 of **FB 1**.
Place the remaining sample of **FB 7** in a boiling tube and add portions of **FB 1** from the measuring cylinder until all the **FB 7** has reacted. Stir until the reaction is complete.
Record your observations.

Keep this solution for use in (b)(iii).

.....

.....

.....

..... [2]

(iii) Carry out the following tests. For each test use a 1 cm depth of the solution from **(b)(ii)** in a test-tube.
Record your observations in Table 3.1.
Identify the metal ion present in **FB 7**.

Table 3.1

test	*observations*
Test 1 Add aqueous sodium hydroxide.	
Test 2 Add aqueous ammonia.	
Test 3 Add a piece of aluminium foil.	

The formula of the metal ion in **FB 7** is .....

[4]

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

(c) **FB 8** is an acidified aqueous solution containing a metal cation which is listed in the Qualitative analysis notes.

(i) Carry out the following tests. For each test use a 1 cm depth of **FB 8** in a test-tube.
Record your observations in Table 3.2.
Identify the metal ion present in **FB 8**.

Table 3.2

test	*observations*
Test 1 Add aqueous sodium hydroxide.	
Test 2 Add a 1 cm depth of acidified aqueous potassium manganate(VII).	

The formula of the metal ion in **FB 8** is .....

[3]

(ii) State the type of reaction between the metal ion in **FB 8** and acidified aqueous potassium manganate(VII).
Explain how your observations support your answer.

.....

.....

..... [1]

[Total: 15]

Qualitative analysis notes

1 Reactions of cations

cation	reaction with	
	NaOH(aq)	**NH_3(aq)**
aluminium, Al^{3+}(aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH_4^+(aq)	no ppt. ammonia produced on warming	–
barium, Ba^{2+}(aq)	faint white ppt. is observed unless [Ba^{2+}(aq)] is very low	no ppt.
calcium, Ca^{2+}(aq)	white ppt. unless [Ca^{2+}(aq)] is very low	no ppt.
chromium(III), Cr^{3+}(aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess
copper(II), Cu^{2+}(aq)	pale blue ppt. insoluble in excess	pale blue ppt. soluble in excess giving dark blue solution
iron(II), Fe^{2+}(aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe^{3+}(aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
magnesium, Mg^{2+}(aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn^{2+}(aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn^{2+}(aq)	white ppt. soluble in excess	white ppt. soluble in excess

2 Reactions of anions

anion	reaction
carbonate, CO_3^{2-}	CO_2 liberated by dilute acids
chloride, Cl^-(aq)	gives white ppt. with Ag^+(aq) (soluble in NH_3(aq))
bromide, Br^-(aq)	gives cream / off-white ppt. with Ag^+(aq) (partially soluble in NH_3(aq))
iodide, I^-(aq)	gives pale yellow ppt. with Ag^+(aq) (insoluble in NH_3(aq))
nitrate, NO_3^-(aq)	NH_3 liberated on heating with OH^-(aq) and Al foil
nitrite, NO_2^-(aq)	NH_3 liberated on heating with OH^-(aq) and Al foil; decolourises acidified aqueous $KMnO_4$
sulfate, SO_4^{2-}(aq)	gives white ppt. with Ba^{2+}(aq) (insoluble in excess dilute strong acids); gives white ppt. with high [Ca^{2+}(aq)]
sulfite, SO_3^{2-}(aq)	gives white ppt. with Ba^{2+}(aq) (soluble in excess dilute strong acids); decolourises acidified aqueous $KMnO_4$
thiosulfate, $S_2O_3^{2-}$(aq)	gives off-white / pale yellow ppt. slowly with H^+

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

3 Tests for gases

gas	test and test result
ammonia, NH_3	turns damp red litmus paper blue
carbon dioxide, CO_2	gives a white ppt. with limewater
hydrogen, H_2	'pops' with a lighted splint
oxygen, O_2	relights a glowing splint

4 Tests for elements

element	test and test result
iodine, I_2	gives blue-black colour on addition of starch solution

Important values, constants and standards

molar gas constant	$R = 8.31\,J\,K^{-1}\,mol^{-1}$
Faraday constant	$F = 9.65 \times 10^4\,C\,mol^{-1}$
Avogadro constant	$L = 6.02 \times 10^{23}\,mol^{-1}$
electronic charge	$e = -1.60 \times 10^{-19}\,C$
molar volume of gas	$V_m = 22.4\,dm^3\,mol^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0\,dm^3\,mol^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14}\,mol^2\,dm^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18\,kJ\,kg^{-1}\,K^{-1}$ ($4.18\,J\,g^{-1}\,K^{-1}$)

© UCLES 2025

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

The Periodic Table of Elements

Group																	
1	2											13	14	15	16	17	18
							1 H hydrogen 1.0										2 He helium 4.0
3 Li lithium 6.9	4 Be beryllium 9.0											5 B boron 10.8	6 C carbon 12.0	7 N nitrogen 14.0	8 O oxygen 16.0	9 F fluorine 19.0	10 Ne neon 20.2
11 Na sodium 23.0	12 Mg magnesium 24.3	3	4	5	6	7	8	9	10	11	12	13 Al aluminium 27.0	14 Si silicon 28.1	15 P phosphorus 31.0	16 S sulfur 32.1	17 Cl chlorine 35.5	18 Ar argon 39.9
19 K potassium 39.1	20 Ca calcium 40.1	21 Sc scandium 45.0	22 Ti titanium 47.9	23 V vanadium 50.9	24 Cr chromium 52.0	25 Mn manganese 54.9	26 Fe iron 55.8	27 Co cobalt 58.9	28 Ni nickel 58.7	29 Cu copper 63.5	30 Zn zinc 65.4	31 Ga gallium 69.7	32 Ge germanium 72.6	33 As arsenic 74.9	34 Se selenium 79.0	35 Br bromine 79.9	36 Kr krypton 83.8
37 Rb rubidium 85.5	38 Sr strontium 87.6	39 Y yttrium 88.9	40 Zr zirconium 91.2	41 Nb niobium 92.9	42 Mo molybdenum 95.9	43 Tc technetium –	44 Ru ruthenium 101.1	45 Rh rhodium 102.9	46 Pd palladium 106.4	47 Ag silver 107.9	48 Cd cadmium 112.4	49 In indium 114.8	50 Sn tin 118.7	51 Sb antimony 121.8	52 Te tellurium 127.6	53 I iodine 126.9	54 Xe xenon 131.3
55 Cs caesium 132.9	56 Ba barium 137.3	57–71 lanthanoids	72 Hf hafnium 178.5	73 Ta tantalum 180.9	74 W tungsten 183.8	75 Re rhenium 186.2	76 Os osmium 190.2	77 Ir iridium 192.2	78 Pt platinum 195.1	79 Au gold 197.0	80 Hg mercury 200.6	81 Tl thallium 204.4	82 Pb lead 207.2	83 Bi bismuth 209.0	84 Po polonium –	85 At astatine –	86 Rn radon –
87 Fr francium –	88 Ra radium –	89–103 actinoids	104 Rf rutherfordium –	105 Db dubnium –	106 Sg seaborgium –	107 Bh bohrium –	108 Hs hassium –	109 Mt meitnerium –	110 Ds darmstadtium –	111 Rg roentgenium –	112 Cn copernicium –	113 Nh nihonium –	114 Fl flerovium –	115 Mc moscovium –	116 Lv livermorium –	117 Ts tennessine –	118 Og oganesson –

Key

atomic number
atomic symbol
name
relative atomic mass

lanthanoids	57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.2	61 Pm promethium –	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0
actinoids	89 Ac actinium –	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium –	94 Pu plutonium –	95 Am americium –	96 Cm curium –	97 Bk berkelium –	98 Cf californium –	99 Es einsteinium –	100 Fm fermium –	101 Md mendelevium –	102 No nobelium –	103 Lr lawrencium –

DO NOT WRITE IN THIS MARGIN DO NOT WRITE IN THIS MARGIN DO NOT WRITE IN THIS MARGIN DO NOT WRITE IN THIS MARGIN DO NOT WRITE IN THIS MARGIN