CAMBRIDGE
International Education

Cambridge International AS & A Level

PHYSICS **9702/54**

Paper 5 Planning, Analysis and Evaluation **October/November 2025**

MARK SCHEME

Maximum Mark: 30

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2025 series for most Cambridge IGCSE, Cambridge International A and AS Level components, and some Cambridge O Level components.

© Cambridge University Press & Assessment 2025

PUBLISHED

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptions for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

© Cambridge University Press & Assessment 2025

GENERIC MARKING PRINCIPLE 5: Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).
GENERIC MARKING PRINCIPLE 6: Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Science-Specific Marking Principles

1	Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.
2	The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.
3	Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).
4	The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.
5	<u>'List rule' guidance</u> For questions that require ***n*** responses (e.g. State **two** reasons …): • The response should be read as continuous prose, even when numbered answer spaces are provided. • Any response marked *ignore* in the mark scheme should not count towards ***n***. • Incorrect responses should not be awarded credit but will still count towards ***n***. • Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should **not** be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response. • Non-contradictory responses after the first ***n*** responses may be ignored even if they include incorrect science.

© Cambridge University Press & Assessment 2025

6 Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form (e.g. $a \times 10^n$) in which the convention of restricting the value of the coefficient (a) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

7 Guidance for chemical equations

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.

© Cambridge University Press & Assessment 2025

Annotations guidance for centres

Examiners use a system of annotations as a shorthand for communicating their marking decisions to one another. Examiners are trained during the standardisation process on how and when to use annotations. The purpose of annotations is to inform the standard isation and monitoring processes and guide the supervising examiners when they are checking the work of examiners within their team. The meaning of annotations and how they are used is specific to each component and is understood by all examiners who mark the component.

We publish annotations in our mark schemes to help centres understand the annotations they may see on copies of scripts. Note that there may not be a direct correlation between the number of annotations on a script and the mark awarded. Similarly, the use of an annotation may not be an indication of the quality of the response.

The annotations listed below were available to examiners marking this component in this series.

Annotations

Annotation	Meaning
BOD	benefit of the doubt given
1	correct awarding one mark from additional detail 1. similar numbered ticks are used for additional detail 2, 3, 4 etc.
✓	correct point or mark awarded
P	defining the problem mark
ECF	error carried forward applied
SF	error in number of significant figures
I	incorrect or insufficient point ignored while marking the rest of the response
✗	incorrect point or mark not awarded
U	incorrect unit
^	information missing or insufficient for credit

© Cambridge University Press & Assessment 2025

Annotation	Meaning
MO	methods of data collection mark
SEEN	point has been noted, but no credit has been given or blank page seen
R	repeat of point previously awarded mark

© Cambridge University Press & Assessment 2025

Question	Answer	Marks
1	**Defining the problem**	
	vary m and measure f **or** m is the independent variable and f is the dependent variable	**1**
	keep f_0 constant	**1**
	Methods of data collection	
	labelled diagram of workable experiment including: • turntable base and motor on bench • two labels from turntable, motor, belt, C, putty, bench, release mechanism	**1**
	workable circuit showing terminals of motor connected to (d.c.) supply	**1**
	method to measure time to determine f and f_0, e.g. use a stopwatch to measure (n)t **and** method to measure m, e.g. use a (top-pan) balance	**1**
	measure the time t for n revolutions **and** $T = t / n$ **and** $f = 1 / T$ **or** $f = n / t$	**1**
	Method of Analysis	
	plots a graph of $\frac{1}{f}$ against m or equivalent Do not accept logarithms.	**1**

© Cambridge University Press & Assessment 2025

Question	Answer	Marks
1	$\frac{1}{f}$ against m | m against $\frac{1}{f}$ $K = \frac{r^2}{f_0 \times \text{gradient}}$ | $K = \frac{r^2 \times \text{gradient}}{f_0}$	**1**
	$\frac{1}{f}$ against m | m against $\frac{1}{f}$ $\beta = f_0 \times y\text{-intercept}$ | $\beta = -\frac{f_0 \times y\text{-intercept}}{\text{gradient}}$ **or** $\beta = -\frac{r^2 \times y\text{-intercept}}{K}$	**1**

Question	Answer	Marks
1	**Additional detail including safety considerations**	**6**
	D1 precaution linked to <u>putty</u> leaving turntable, e.g. screens around apparatus / cushions to prevent putty from leaving bench	
	D2 keep *r* <u>constant</u>	
	D3 use rule(r) to measure *r*	
	D4 stand and fixed point located above P (to ensure *r* is constant)	
	D5 method to ensure putty is dropped at (constant) distance *r*, e.g.; fixed location above the turntable to drop putty to keep *r* constant **or** draw a circle of radius *r* on the turntable	
	D6 method to keep f_0 constant, e.g. adjust the rheostat to keep the current in the motor constant / check ammeter	
	D7 clamp motor and/or turntable (base) <u>to bench</u>	
	D8 lubricate the turntable to reduce friction	
	D9 repeat experiment for the same value of *m* **and** determine the average *f*	
	D10 relationship valid <u>if</u> a straight line is produced (with *y*-intercept = $\frac{\beta}{f_0}$) Do not accept line through the origin.	

© Cambridge University Press & Assessment 2025

<table>
<tr><th>Question</th><th>Answer</th><th>Marks</th></tr>
<tr><td>2(a)</td><td>gradient = $D\lambda$</td><td>1</td></tr>
<tr><td rowspan="2">2(b)</td><td>
<table>
<tr><th>$\frac{1}{s}$ / mm^{-1}</th><th>y / mm</th></tr>
<tr><td>5.6 or 5.56</td><td>3.30</td></tr>
<tr><td>4.8 or 4.76</td><td>2.89</td></tr>
<tr><td>4.2 or 4.17</td><td>2.51</td></tr>
<tr><td>3.7 or 3.70</td><td>2.26</td></tr>
<tr><td>3.2 or 3.23</td><td>1.96</td></tr>
<tr><td>2.6 or 2.63</td><td>1.59</td></tr>
</table>
Values of $\frac{1}{s}$ and y correct as shown above.</td><td>1</td></tr>
<tr><td>Uncertainties in $\frac{1}{s}$ from $\pm$ 0.3 to $\pm$ 0.07.</td><td>1</td></tr>
<tr><td rowspan="2">2(c)(i)</td><td>Six points from (b) plotted correctly.
Must be within half a small square. Diameter of points must be less than half a small square.</td><td>1</td></tr>
<tr><td>Error bars in $\frac{1}{s}$ plotted correctly.
All error bars must be plotted. Total length of bar must be accurate to less than half a small square and symmetrical.</td><td>1</td></tr>
</table>

© Cambridge University Press & Assessment 2025

Question	Answer	Marks
2(c)(ii)	Straight line of best fit drawn. Thickness of the line must be less than half a small square. Do not accept line from top point to bottom point. Line must pass between (2.90, 1.80) and (3.00, 1.80) **and** between (5.30, 3.20) and (5.40, 3.20).	**1**
	Worst acceptable line drawn (steepest or shallowest possible line that passes through all the error bars). Thickness of the line must be less than half a small square. All error bars must be plotted.	**1**
2(c)(iii)	Gradient determined with clear substitution of data points into $\Delta y / \Delta x$. Distance between data points must be greater than half the length of the drawn line.	**1**
	Gradient determined of worst acceptable line with clear substitution of data points into $\Delta y / \Delta x$. uncertainty = (gradient of line of best fit – gradient of worst acceptable line) **or** uncertainty = ½ (steepest worst line gradient – shallowest worst line gradient)	**1**
2(d)	(0.921 ± 0.008) (m)	**1**
2(e)(i)	λ determined using gradient **and** λ given to 2 or 3 significant figures. $\lambda = \frac{\text{gradient}}{D}$	**1**
	λ determined using gradient **and** λ given with SI units with appropriate power of ten. $\lambda = \frac{\text{gradient}}{D}$ Unit of λ: m or mm.	**1**

© Cambridge University Press & Assessment 2025

Question	Answer	Marks
2(e)(ii)	Percentage uncertainty in λ determined with clear method shown. $\%\text{uncertainty} = \left(\frac{\Delta\text{gradient}}{\text{gradient}} + \frac{\Delta D}{D}\right) \times 100$ **or** correct substitution for max/min methods.	**1**
2(f)	s determined to a minimum of 2 significant figures from **(c)(iii)** or **(d)** and **(e)(i)** with correct substitution **and** correct power of ten. $s = \frac{\text{gradient}}{0.005\ (\text{m})}$ **or** $s = \frac{D \times \lambda}{0.005\ (\text{m})} = \frac{\textbf{(d)} \times \textbf{(e)(i)}}{0.005\ (\text{m})}$	**1**
	Absolute uncertainty determined with correct substitution. For use of gradient from **(c)(iii)**: $\Delta s = \left(\frac{\Delta\text{gradient}}{\text{gradient}} + \frac{0.005}{0.500}\right) \times s$ **or** correct substitution for max/min methods. For use of D and λ from **(d)** and **(e)(i)**: $\Delta s = \left(\frac{\Delta D}{D} + \frac{\Delta\lambda}{\lambda} + \frac{0.005}{0.500}\right) \times s$ **or** correct substitution for max/min methods: $\Delta s = \left(\frac{\max D \times \max\lambda}{\min y}\right) - s$ **or** $\Delta s = s - \left(\frac{\min D \times \min\lambda}{\max y}\right)$	**1**

© Cambridge University Press & Assessment 2025