Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER			

9337495756

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **February/March 2023**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages.

© UCLES 2023

1 It is given that $x = \ln(2y - 3) - \ln(y + 4)$.

Express y in terms of x. [3]

2 **(a)** On an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $-\frac{1}{3}\pi \leqslant \arg(z - 1 - 2\mathrm{i}) \leqslant \frac{1}{3}\pi$ and $\mathrm{Re}\, z \leqslant 3$. [3]

(b) Calculate the least value of $\arg z$ for points in the region from **(a)**. Give your answer in radians correct to 3 decimal places. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3 The polynomial $2x^4 + ax^3 + bx - 1$, where a and b are constants, is denoted by $\mathrm{p}(x)$. When $\mathrm{p}(x)$ is divided by $x^2 - x + 1$ the remainder is $3x + 2$.

Find the values of a and b. [5]

4 Solve the equation

$$\frac{5z}{1+2\mathrm{i}} - zz^* + 30 + 10\mathrm{i} = 0,$$

giving your answers in the form $x + \mathrm{i}y$, where x and y are real. [5]

5 The parametric equations of a curve are

$$x = te^{2t}, \qquad y = t^2 + t + 3.$$

(a) Show that $\frac{dy}{dx} = e^{-2t}$. [3]


(b) Hence show that the normal to the curve, where $t = -1$, passes through the point $\left(0,\ 3 - \frac{1}{e^4}\right)$. [3]

6 **(a)** Express $5\sin\theta + 12\cos\theta$ in the form $R\cos(\theta - \alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$. [3]

© UCLES 2023

(b) Hence solve the equation $5 \sin 2x + 12 \cos 2x = 6$ for $0 \leqslant x \leqslant \pi$. [4]

7



The diagram shows a circle with centre O and radius r. The angle of the **minor** sector AOB of the circle is x radians. The area of the **major** sector of the circle is 3 times the area of the shaded region.

(a) Show that $x = \frac{3}{4}\sin x + \frac{1}{2}\pi$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Show by calculation that the root of the equation in **(a)** lies between 2 and 2.5. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use an iterative formula based on the equation in **(a)** to calculate this root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

8



The diagram shows the curve $y = x^3 \ln x$, for $x > 0$, and its minimum point M.

(a) Find the exact coordinates of M. [4]

(b) Find the exact area of the shaded region bounded by the curve, the x-axis and the line $x = \frac{1}{2}$. [5]

9 The variables x and y satisfy the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \mathrm{e}^{3y} \sin^2 2x.$$

It is given that $y = 0$ when $x = 0$.

Solve the differential equation and find the value of y when $x = \frac{1}{2}$. [7]

10 With respect to the origin O, the points A, B, C and D have position vectors given by

$$\overrightarrow{OA} = \begin{pmatrix} 3 \\ -1 \\ 2 \end{pmatrix}, \quad \overrightarrow{OB} = \begin{pmatrix} 1 \\ 2 \\ -3 \end{pmatrix}, \quad \overrightarrow{OC} = \begin{pmatrix} 1 \\ -2 \\ 5 \end{pmatrix} \quad \text{and} \quad \overrightarrow{OD} = \begin{pmatrix} 5 \\ -6 \\ 11 \end{pmatrix}.$$

(a) Find the obtuse angle between the vectors $\overrightarrow{OA}$ and $\overrightarrow{OB}$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The line l passes through the points A and B.

(b) Find a vector equation for the line l. [2]

.....

.....

.....

.....

.....

.....

(c) Find the position vector of the point of intersection of the line l and the line passing through C and D. [4]

11 Let $\mathrm{f}(x) = \dfrac{5x^2 + x + 11}{(4 + x^2)(1 + x)}$.

(a) Express $\mathrm{f}(x)$ in partial fractions. [5]

(b) Hence show that $\int_0^2 \mathrm{f}(x)\,\mathrm{d}x = \ln 54 - \frac{1}{8}\pi$. [5]

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2023