**CAMBRIDGE INTERNATIONAL EXAMINATIONS**

**GCE Advanced Subsidiary Level and GCE Advanced Level**

# MARK SCHEME for the May/June 2013 series

## 9709 MATHEMATICS

**9709/61** Paper 6, maximum raw mark 50

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

CAMBRIDGE
International Examinations

**Mark Scheme Notes**

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.

- The symbol √ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.

- Note: B2 or A2 means that the candidate can earn 2 or 0.
  B2/1/0 means that the candidate can earn anything from 0 to 2.

  The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.

- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking *g* equal to 9.8 or 9.81 instead of 10.

© Cambridge International Examinations 2013

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF Any Equivalent Form (of answer is equally acceptable)

AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)

BOD Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)

CAO Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)

CWO Correct Working Only – often written by a 'fortuitous' answer

ISW Ignore Subsequent Working

MR Misread

PA Premature Approximation (resulting in basically correct work that is insufficiently accurate)

SOS See Other Solution (the candidate makes a better attempt at the same question)

SR Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

**Penalties**


MR –1 A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through √" marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR –2 penalty may be applied in particular cases if agreed at the coordination meeting.

PA –1 This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

© Cambridge International Examinations 2013

| | | | | | |
|---|---|---|---|---|---|
| **1** | **(i)** | $sd^2 = 1957.5/30 - (234/30)^2$<br>$sd = 2.1$ | M1<br>A1 | **[2]** | Subst in formula or expand<br>Accept 2.10 |
| | **(ii)** | $86 = 234/30 + c$<br>$c = 78.2$ | M1<br>A1 | **[2]** | 234/30 seen |
| **2** | | $np = 350 \times 1/7$ (= 50)<br>$npq = 350 \times 1/7 \times 6/7$ (= 42.857)<br>$P(x \quad 47) = P\left(z > \frac{46.5-50}{\sqrt{42.857}}\right) =$<br>$P(z > -0.5346)$<br>= 0.704 | B1<br>M1<br>M1<br>M1<br>A1 | **[5]** | Correct unsimplified *np* and *npq*<br>standardising, with or without cc, must have sq rt<br>continuity correction 46.5 or 47.5<br>correct area ie > 0.5 must be a Φ<br>correct answer |
| **3** | **(i)** | females: med \$22 700<br>LQ \$21700 UQ \$24 000 | B1<br>B1 | **[2]** | Any 2 correct<br>All correct |
| | **(ii)** | males<br>females<br>20 21 22 23 24 25 26 27<br>Salary in \$000 | B1<br>B1<br>B1 | **[3]** | Uniform scale and labels must see Salary, \$000<br>Correct graph for females ft their quartiles. Line not through box<br>Correct graph for males |
| **4** | **(a)** | $P(y < 0) = P\left(z < \frac{0-\mu}{\mu/2}\right)$<br>$= P(z < -2)$<br>$= 1 - 0.9772 = 0.0228$ | M1<br>A1<br>A1 | **[3]** | Standardising containing 0 (can be implied) and μ only<br>$z < -2$ seen<br>Correct answer |
| **(b)** | | $P(x > 2.1) = 253/8000 = 0.031625$<br>$P(x < 2.1) = 0.968375 = \Phi(z)$<br>$z = 1.857$ or 1.858 or 1.859 $= \frac{2.1-2.04}{\sigma}$<br>$\sigma = 0.0323$ | M1<br>A1<br>M1<br>A1 | **[4]** | 1 – their 253/8000 used to obtain a *z*-value<br>Rounded to 1.86 seen<br>Solving for σ using their *z* val must be a *z* val<br>Correct answer |
| **5** | **(i)** | $X \sim$ Bin (12, 0.2) | B1<br>B1<br>B1 | **[3]** | Bin or B<br>12<br>0.2 or 1/5 |
| | **(ii)** | $P(X = 3, 4, 5) = 0.2^3 0.8^9 {}_{12}C_3 + 0.2^4 0.8^8 {}_{12}C_4 + 0.2^5 0.8^7 {}_{12}C_5$<br>= 0.23622 + 0.13287 + 0.05315<br>= 0.422 | M1<br>A1ft<br>A1 | **[3]** | Bin exprerssion with any p<br>Correct unsimplified expression, their p<br>Correct answer |

© Cambridge International Examinations 2013

| | | | | | |
|---|---|---|---|---|---|
| | **(iii)** | P ($X = 0$) < 0.01<br><br>$0.8^n < 0.01$<br>$n = 21$ | M1<br><br>M1<br>A1 | **[3]** | Statement involving P(X = 0) and 0.01 can be implied<br>Equn involving '0.8', 0.01 or 0.99<br>Correct answer |
| **6** | **(i)** | 4! × 3! × 5! × 2! × 4! = 829440 | B1<br><br>B1<br>B1 | **[3]** | 4!, 3!, 5!, 2 seen multiplied 1, not in denominator<br>Mult by 4!<br>Correct answer |
| | **(ii)** | 8! × 9 × 8 × 7 × 6 × 5 × 4<br><br>= 2438553600 ($2.44 \times 10^9$) | B1<br>B1<br>B1 | **[3]** | 8! seen multiplied 1<br>Mult by ${}_9P_6$<br>Correct answer |
| | **(iii)** | 8C3 × 5C3 × 2C2<br>= 560 | B1<br>B1<br>B1 | **[3]** | 8C3 seen mult<br>5C3 seen mult<br>Correct answer |
| **7** | **(i)** | number of balls in $B$ is $5 + x + 1 = x + 6$<br>P($Y$) = $x/(x + 6)$ **AG** | B1 | **[1]** | Sensible reason |
| | **(ii)** | box $A$ box $B$<br><br> | B1<br><br><br><br>B1<br><br>B1<br><br><br>B1 | **[4]** | both correct for box A<br><br><br><br>1 correct<br><br>1 correct<br><br><br>1 correct |
| | **(iii)** | P($W_B$) = $\frac{6}{x+6} = \frac{1}{3}$<br><br>$x = 12$ **AG** | M1<br><br>A1 | **[2]** | their $\frac{6}{x+6}$ = 1/3 or $x/x$+6 = 2/3<br><br>Verification or solving legit |

© Cambridge International Examinations 2013

| | | | | |
|---|---|---|---|---|
| **(iv)** | $P(Y) = \frac{8}{10} \times \frac{12}{18} + \frac{2}{10} \times \frac{13}{18}$ | M1 | | Attempt at P(,$Y$) involving 2 two-factor fractions, seen anywhere. |
| | $= \frac{61}{90}$ | A1 | | Correct P(Y) seen as num or denom of a fraction |
| | $P( = (AY \mid BY) = \frac{P(AY \cap BY)}{P(Y)}$ | B1 | | (2/10) × (13/18) seen as num or denom of a fraction |
| | $= \frac{2}{10} \times \frac{13}{18} / \frac{61}{90}$ | | | |
| | $= \frac{13}{61} (0.213)$ | A1 | **[4]** | Correct answer |

© Cambridge International Examinations 2013