CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Level

MARK SCHEME for the May/June 2014 series

9709 MATHEMATICS

9709/72 Paper 7, maximum raw mark 50

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2014 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.

- The symbol √ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.

- Note: B2 or A2 means that the candidate can earn 2 or 0.
  B2/1/0 means that the candidate can earn anything from 0 to 2.

  The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.

- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

© Cambridge International Examinations 2014

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF Any Equivalent Form (of answer is equally acceptable)

AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)

BOD Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)

CAO Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)

CWO Correct Working Only – often written by a 'fortuitous' answer

ISW Ignore Subsequent Working

MR Misread

PA Premature Approximation (resulting in basically correct work that is insufficiently accurate)

SOS See Other Solution (the candidate makes a better attempt at the same question)

SR Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

MR –1 A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through √" marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR –2 penalty may be applied in particular cases if agreed at the coordination meeting.

PA –1 This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

© Cambridge International Examinations 2014

1	$\frac{\Sigma x}{8} = \frac{2006}{8} = 250.75$ or 251 (3 s.f.) $(\Sigma x^2 = 503274)$ $\frac{8}{7}\left(\frac{"503274"}{8} - "250.75"^2\right)$ = 38.5 o.e. (accept 6.204²)	B1 M1 A1 [3]	Any equivalent form For use of formula of correct form cao (as final answer)
2	$(X + Y - Z) \sim N(8, \ldots)$ μ=8 (or –8) Var$(X + Y - Z) = 2^2 + 1.5^2 + 1.8^2$ (= 9.49) $\frac{0-8}{\sqrt{'9.49'}}$ (= –2.597) Φ('–2.597') = 1 – Φ('2.597') = 0.0047	 B1 B1 M1 M1 A1 [5]	 seen or implied – award at early stage For standardising (accept sd/var mixes, but variance must be a combination of at least 2 of X, Y, Z) For area consistent with their working
3	H_0: Pop mean (or μ or λ) = 50 (or 5) H_1: Pop mean (or μ or λ) ≠ 50 (or 5) $\frac{60.5-50}{\sqrt{50}}$ (±) = (±)1.485 OR 0.0687 OR C.V 1.485 < 1.645 or 0.0687 > 0.05 No evidence that mean changed	 B1 M1 A1 M1 A1√ [5]	 Not just "mean" For standardising with N(50,50) or N(5,5/√10) Allow M1 with wrong or no continuity correction OR no √ (accept c.v method M1, A1 for 61.63 or 48.868) For valid comparison (zs or areas or cv) (S.R For cv comparison 61.63 only award final A1 if cc used) or if H_1: $\lambda > 50$, 1.485 < 1.96 M1 No evid mean changed A0 (i.e. if one-tail test, max B0 M1 A1 M1 A0)

© Cambridge International Examinations 2014

4	**(i)**	$\lambda = 4.5$	B1	seen
		$1 - e^{-4.5}\left(1+4.5+\frac{4.5^2}{2}\right)$	M1	any λ. Allow one end error
		$= 0.826$ (3 s.f.)	A1 [3]	
	(ii)	$e^{-\lambda} = 0.523$	B1	
		$(-\lambda = \ln 0.523)$		
		$\lambda = 0.648$ (3 s.f.)	B1 [2]	
	(iii)	$e^{-\mu} \times \frac{\mu^3}{3!} = 24 \times e^{-\mu} \times \mu$	B1	
		$\frac{\mu^2}{6} = 24$	M1	For a simplified expression in μ^2 with $e^{-\mu}$ and μ cancelled and no factorials.
		$\mu = 12$	A1 [3]	
5	**(i)**	$p = \frac{184}{400}$ or 0.46	B1	Used
		$z = 1.96$	B1	Seen
		$"0.46" \pm z \times \sqrt{\frac{"0.46"(1-"0.46")}{400}}$	M1	Using expression of correct form
		$= 0.411$ to 0.509	A1 [4]	Must be an interval
	(ii)	0.5 within CI		
		Claim not supported or not justified	B1√ [1]	Both needed. No contradictions. ft their **(i)**
	(iii)	$z \times \sqrt{\frac{"0.46"(1-"0.46")}{400}} = 0.05$	M1	Allow M1 for $z \times \sqrt{\frac{"0.46"(1-"0.46")}{400}} = 0.1$
		$z = 2.006$	A1	
		$\Phi(\text{'2.006'}) = 0.9775$		
		$\alpha = \text{'0.9775'} - (1 - \text{'0.9775'})$	M1	or $1 - 2(1 - \text{'0.9775'})$
		$= 95.5\%$	A1 [4]	

© Cambridge International Examinations 2014

6 (i)	$k\int_0^4(16t-t^3)\mathrm{d}t = 1$	M1	Int f(t) = 1 ignore limits
	$k\left[8t^2-\frac{t^4}{4}\right]_0^4 = 1$	A1	correct integration with correct limits
	$k(128-64) = 1$ o.e.		
	$k \times 64 = 1$		
	$\left(k=\frac{1}{64}\right)$ **AG**	A1 [3]	must be convinced (AG)
(ii)	$\frac{1}{64}\int_0^1(16t-t^3)\mathrm{d}t$	M1	Int f(t) between 0 and 1 (accept 0 and a value < 1 , 1 and 4)
	$= \frac{1}{64}\left[8t^2-\frac{t^4}{4}\right]_0^1$	A1	correct integration and correct limits (ignore "k")
	$= \frac{1}{64}\left[8-\frac{1}{4}\right]$		
	$= \frac{31}{256}$ or 0.121094	A1	
	$\left(\frac{31}{256}\right)^2 = 0.0147$ (3 s.f.) o.e.	B1√ [4]	ft their "$\frac{31}{256}$"
iii	$\frac{1}{64}\int_0^4(16t^2-t^4)\mathrm{d}t$	M1	Int tf(t) ignore limits
	$= \frac{1}{64}\left[\frac{16t^3}{3}-\frac{t^5}{5}\right]_0^4$	A1	correct integration and correct limits (ignore "k")
	$= \frac{1}{64}\left(\frac{1024}{3}-\frac{1024}{5}\right)$		
	$= \frac{32}{15}$ or 2.13 (3 s.f.) o.e.	A1 [3]	

© Cambridge International Examinations 2014

7 (i)	2^{nd} More representative of all appointments or Lengths may vary during the day or 1^{st} does not include later appts so not representative	B1 B1 [2]	Any implication that times or conditions vary throughout day, e.g. doctors get tired	
(ii)	0.01 o.e. Concluding that times spent are too long when they are not.	B1 B1 [2]	Concluding that the mean time spent is more than 10 mins when it is not. Must be in context.	
(iii)	H_0: Pop mean appt time (or μ) = 10 H_1: Pop mean appt time (or μ) > 10 $\frac{\frac{147}{12}-10}{\frac{3.4}{\sqrt{12}}}$ (±) = (±)2.292 or (0.0109 if area comparison done) "2.292" < 2.326 o.e. (No evidence to reject H_0.) No reason to believe appts are too long	B1 M1 A1 M1 A1√ [5]	Both correct. Allow μ, but not just "mean" Allow incorrect $\frac{147}{12}$ Must have √12 (accept totals method) For valid comparison Comp "2.292" with 2.326 Or 0.0109 with 0.01 Or 147/12 with 12.28 Dep 2.326, ft their "2.292" No contradictions	$10 + 2.326 \times \frac{3.4}{\sqrt{12}}$ M1 = 12.28 A1 $\frac{147}{12} < 12.28$ M1
(iv)	Normal population	B1 [1]	Must have "population" or equiv	

© Cambridge International Examinations 2014