Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

5471210840

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **May/June 2023**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages.

© UCLES 2023

1 Solve the equation $\ln(x + 5) = 5 + \ln x$. Give your answer correct to 3 decimal places. [4]

2 Find the quotient and remainder when $2x^4 - 27$ is divided by $x^2 + x + 3$. [3]

3 On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - 3 - \mathrm{i}| \leqslant 3$ and $|z| \geqslant |z - 4\mathrm{i}|$. [4]


© UCLES 2023

4 The parametric equations of a curve are

$$x = \frac{\cos\theta}{2 - \sin\theta}, \qquad y = \theta + 2\cos\theta.$$

Show that $\frac{\mathrm{d}y}{\mathrm{d}x} = (2 - \sin\theta)^2$. [5]

5



The diagram shows the part of the curve $y = x^2 \cos 3x$ for $0 \leqslant x \leqslant \frac{1}{6}\pi$, and its maximum point M, where $x = a$.

(a) Show that a satisfies the equation $a = \frac{1}{3}\tan^{-1}\left(\frac{2}{3a}\right)$. [3]

(b) Use an iterative formula based on the equation in **(a)** to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

6 **(a)** Express $3\cos x + 2\cos(x - 60°)$ in the form $R\cos(x - \alpha)$, where $R > 0$ and $0° < \alpha < 90°$. State the exact value of R and give α correct to 2 decimal places. [4]

(b) Hence solve the equation

$$3\cos 2\theta + 2\cos(2\theta - 60°) = 2.5$$

for $0° < \theta < 180°$. [4]

7 **(a)** Use the substitution $u = \cos x$ to show that

$$\int_0^{\pi} \sin 2x \, e^{2\cos x} \, dx = \int_{-1}^{1} 2ue^{2u} \, du. \quad [4]$$

(b) Hence find the exact value of $\int_0^{\pi} \sin 2x \, e^{2\cos x} \, dx$. [4]

8 The variables x and y satisfy the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{y^2 + 4}{x(y + 4)}$$

for $x > 0$. It is given that $x = 4$ when $y = 2\sqrt{3}$.

Solve the differential equation to obtain the value of x when $y = 2$. [8]

9 The lines l and m have equations

$$l: \quad \mathbf{r} = a\mathbf{i} + 3\mathbf{j} + b\mathbf{k} + \lambda(c\mathbf{i} - 2\mathbf{j} + 4\mathbf{k}),$$
$$m: \quad \mathbf{r} = \mathbf{i} + 2\mathbf{j} + 3\mathbf{k} + \mu(2\mathbf{i} - 3\mathbf{j} + \mathbf{k}).$$

Relative to the origin O, the position vector of the point P is $4\mathbf{i} + 7\mathbf{j} - 2\mathbf{k}$.

(a) Given that l is perpendicular to m and that P lies on l, find the values of the constants a, b and c. [4]

(b) The perpendicular from P meets line m at Q. The point R lies on PQ extended, with $PQ : QR = 2 : 3$.

Find the position vector of R. [6]

10 Let $f(x) = \dfrac{21 - 8x - 2x^2}{(1 + 2x)(3 - x)^2}$.

(a) Express $f(x)$ in partial fractions. [5]

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [5]

11 The complex number z is defined by $z = \dfrac{5a - 2\mathrm{i}}{3 + a\mathrm{i}}$, where a is an integer. It is given that $\arg z = -\frac{1}{4}\pi$.

(a) Find the value of a and hence express z in the form $x + \mathrm{i}y$, where x and y are real. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Express z^3 in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leqslant \pi$. Give the simplified exact values of r and θ. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2023