UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education Advanced Subsidiary Level

MATHEMATICS **9709/22**

Paper 2 Pure Mathematics 2 **(P2)** **October/November 2010**

1 hour 15 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

* 8 6 2 7 7 1 2 7 2 3 *

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use a soft pencil for any diagrams or graphs.
Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 50.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

© UCLES 2010

1 Solve the inequality $|x + 1| > |x - 4|$. [3]

2 Use logarithms to solve the equation $5^x = 2^{2x+1}$, giving your answer correct to 3 significant figures. [4]

3 Show that $\int_0^1 (e^x + 1)^2 \, dx = \frac{1}{2}e^2 + 2e - \frac{3}{2}$. [5]

4 The parametric equations of a curve are

$$x = 1 + \ln(t - 2), \quad y = t + \frac{9}{t}, \quad \text{for } t > 2.$$

(i) Show that $\frac{dy}{dx} = \frac{(t^2 - 9)(t - 2)}{t^2}$. [3]

(ii) Find the coordinates of the only point on the curve at which the gradient is equal to 0. [3]

5 Solve the equation $8 + \cot\theta = 2\,\text{cosec}^2\theta$, giving all solutions in the interval $0° \leqslant \theta \leqslant 360°$. [6]

6 The curve with equation $y = \frac{6}{x^2}$ intersects the line $y = x + 1$ at the point P.

(i) Verify by calculation that the x-coordinate of P lies between 1.4 and 1.6. [2]

(ii) Show that the x-coordinate of P satisfies the equation

$$x = \sqrt{\left(\frac{6}{x + 1}\right)}.$$ [2]

(iii) Use the iterative formula

$$x_{n+1} = \sqrt{\left(\frac{6}{x_n + 1}\right)},$$

with initial value $x_1 = 1.5$, to determine the x-coordinate of P correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]


7 The polynomial $3x^3 + 2x^2 + ax + b$, where a and b are constants, is denoted by $p(x)$. It is given that $(x - 1)$ is a factor of $p(x)$, and that when $p(x)$ is divided by $(x - 2)$ the remainder is 10.

(i) Find the values of a and b. [5]

(ii) When a and b have these values, solve the equation $p(x) = 0$. [4]

© UCLES 2010

8



The diagram shows the curve $y = x \sin x$, for $0 \leqslant x \leqslant \pi$. The point $Q\left(\frac{1}{2}\pi, \frac{1}{2}\pi\right)$ lies on the curve.

(i) Show that the normal to the curve at Q passes through the point $(\pi, 0)$. [5]

(ii) Find $\frac{\mathrm{d}}{\mathrm{d}x}(\sin x - x\cos x)$. [2]

(iii) Hence evaluate $\int_0^{\frac{1}{2}\pi} x \sin x \, \mathrm{d}x$. [3]

© UCLES 2010

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.