CAMBRIDGE
International Education

Cambridge International AS & A Level

MATHEMATICS **9709/11**

Paper 1 Pure Mathematics 1 **October/November 2025**

MARK SCHEME

Maximum Mark: 75

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2025 series for most Cambridge IGCSE, Cambridge International A and AS Level components, and some Cambridge O Level components.

© Cambridge University Press & Assessment 2025

PUBLISHED

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptions for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:
Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:
Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:
Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:
Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:
Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:
Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

© Cambridge University Press & Assessment 2025

Mathematics-Specific Marking Principles

1 Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.

2 Unless specified in the question, non-integer answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.

3 Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.

4 Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).

5 Where a candidate has misread a number or sign in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 A or B mark for the misread.

6 Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

© Cambridge University Press & Assessment 2025

Annotations guidance for centres

Examiners use a system of annotations as a shorthand for communicating their marking decisions to one another. Examiners are trained during the standardisation process on how and when to use annotations. The purpose of annotations is to inform the standardisation and monitoring processes and guide the supervising examiners when they are checking the work of examiners within their team. The meaning of annotations and how they are used is specific to each component and is understood by all examiners who mark the component.

We publish annotations in our mark schemes to help centres understand the annotations they may see on copies of scripts. Note that there may not be a direct correlation between the number of annotations on a script and the mark awarded. Similarly, the use of an annotation may not be an indication of the quality of the response.

The annotations listed below were available to examiners marking this component in this series.

Annotations

Annotation	Meaning
	More information required
	Accuracy mark awarded zero
	Accuracy mark awarded one
	Independent accuracy mark awarded zero
	Independent accuracy mark awarded one
	Independent accuracy mark awarded two
	Benefit of the doubt
	Blank Page
	Incorrect
Dep	Used to indicate DM0 or DM1

© Cambridge University Press & Assessment 2025

Annotation	Meaning
DM1	Dependent on the previous M1 mark(s)
FT	Follow through
	Indicate working that is right or wrong
Highlighter	Highlight a key point in the working
ISW	Ignore subsequent work
J	Judgement
JU	Judgement
M0	Method mark awarded zero
M1	Method mark awarded one
M2	Method mark awarded two
MR	Misread
O	Omission or Other solution
Off-page comment	Allows comments to be entered at the bottom of the RM marking window and then displayed when the associated question item is navigated to.
On-page comment	Allows comments to be entered in speech bubbles on the candidate response.
PE	Judgment made by the PE
Pre	Premature approximation
SC	Special case
SEEN	Indicates that work/page has been seen

© Cambridge University Press & Assessment 2025

Annotation	Meaning
SF	Error in number of significant figures
	Correct
TE	Transcription error
XP	Correct answer from incorrect working

© Cambridge University Press & Assessment 2025

PUBLISHED

Mark Scheme Notes

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

DM or **DB** When a part of a question has two or more 'method' steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.

FT Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.

- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
- For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
- The total number of marks available for each question is shown at the bottom of the Marks column.
- Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
- Square brackets [ ] around text or numbers show extra information not needed for the mark to be awarded.

© Cambridge University Press & Assessment 2025

Abbreviations

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no 'follow through' from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

© Cambridge University Press & Assessment 2025

Question	Answer	Marks	Guidance
1	Use of $b^2 - 4ac$	***M1**	
	Obtain $k^2 - 20k + 64$	**A1**	
	Attempt solution of quadratic equation or inequality	**DM1**	Solve quadratic using suitable method. E.g., factorisation, quadratic formula, completing the square.
	Obtain $k < 4,\ k > 16$ or clear equivalent	**B1**	B0 for use of $\leqslant$ and/or $\geqslant$.
		4	

Question	Answer	Marks	Guidance
2(a)	State or imply that second and fifth terms are $a\cos\theta$ and $a\cos^4\theta$ or $r^3 = \frac{1}{64}$	**B1**	Condone poor notation.
	Attempt solution of equation of form $\cos^3\theta = k$ as far as $\cos\theta = \ldots$, where $0 < k < 1$	**M1**	Must come from division.
	Obtain $\cos\theta = 0.25$, and hence $\theta = 1.32$ only	**A1**	Allow greater accuracy 1.318116… A0 for 75.5°.
		3	
2(b)	Attempt to find a using *their* θ and substitute in correct formula for sum to infinity	**M1**	Using exact or approximate value for r. $-1 < r < 1$
	Obtain $a = 32$ and use exact value of r to obtain $\frac{128}{3}$	**A1**	Or exact equivalent.
		2	

© Cambridge University Press & Assessment 2025

Question	Answer	Marks	Guidance
3	x^4 term from $(px+3)^5$ is $5p^4x^4\times 3$	**B1**	Or coefficient is $15p^4$. Not 5C1 × 3 or factorial form or $15(px)^4$. Can be in a list.
	Identify term involving $x^6\times\frac{p^2}{x^2}$ as relevant term from $\left(x^3+\frac{p}{x}\right)^4$	**M1**	
	Obtain $6p^2x^4$ or $-6p^2x^4$	**A1**	Or coefficient is $6p^2$ or $-6p^2$. Not 4C2 or factorial form.
	Attempt solution for p^2 of equation that is quadratic in p^2	**M1**	Solve quadratic in p^2 using suitable method. E.g. factorisation, quadratic formula, completing the square. E.g. $3(5p^2 + 18)(p^2 – 4)$
	Obtain $15p^4-6p^2-216=0$ or equivalent and hence $p=2$	**B1**	No other solutions.
		5	

© Cambridge University Press & Assessment 2025

Question	Answer	Marks	Guidance
4(a)	{10} – (x {+} {3})^2	**B1**	
		B1	
		B1	
		3	
4(b)	State reflection in x-axis	**B1**	
	Obtain $m = -3$	**B1 FT**	Following *their* value of b.
	Obtain $n = 10$	**B1 FT**	Following *their* value of a.
		3	

© Cambridge University Press & Assessment 2025

Question	Answer	Marks	Guidance
5(a)	Use $\tan^4\theta = \frac{\sin^4\theta}{\cos^4\theta}$ or $\tan^4\theta - 1 = \left(\frac{\sin^2\theta}{\cos^2\theta}+1\right)+\left(\frac{\sin^2\theta}{\cos^2\theta}-1\right)$	**B1**	Candidates can work from LHS to RHS or RHS to LHS for full marks. If working on both sides simultaneously, maximum of B1 M1 only if a common correct expression is reached.
	Attempt to express $\tan^4\theta - 1$ as a single fraction in terms of $\cos\theta$ only	**M1**	Using $\frac{(1-c^2)^2-c^4}{c^4}$ or $\frac{(s^2+c^2)(s^2-c^2)}{c^4}$ etc.
	Confirm result $\frac{1-2\cos^2\theta}{\cos^4\theta}$	**A1**	AG – necessary detail needed. WWW
	Alternative Method for Question 5(a)		
	$\frac{1-2\cos^2\theta}{\cos^4\theta} + \frac{\cos^4\theta}{\cos^4\theta} - 1$	**B1**	
	$\frac{\left(1-\cos^2\theta\right)^2}{\cos^4\theta} - 1$	**M1**	
	Confirm result $\tan^4\theta - 1$	**A1**	AG – necessary detail needed.
	Alternative Method 2 for Question 5(a)		
	$\tan^4\theta = \sec^4\theta - 2\sec^2\theta + 1$	**B1**	
	$\tan^4\theta - 1 = \frac{1}{\cos^4\theta} - \frac{2}{\cos^2\theta}$	**M1**	
	Confirm result $\frac{1-2\cos^2\theta}{\cos^4\theta}$	**A1**	AG – necessary detail needed.
		3	

© Cambridge University Press & Assessment 2025

Question	Answer	Marks	Guidance
5(b)	Use identity to form equation using $\cos\theta$ only	***M1**	Expect $\frac{1-2\cos^2\theta}{\cos^2\theta} = 7$ OE Allow unsimplified.
	Obtain $9\cos^2\theta = 1$ or similarly simplified equivalent	**A1**	Condone $\cos^2\theta\ (1 - 9\cos^2\theta) = 0$ Could be implied by one correct solution
	Solve equation of form $\cos^2\theta = k$ to obtain at least one solution	**DM1**	Provided $0 < k < 1$.
	Obtain 70.5 and 109.5	**A1**	Or greater accuracy 70.528…, 109.471…, and no others between 0 and 180. **SC B1** for 1.23^c and 1.91^c only. A0 if $\cos^2\theta\,(1 - 9\cos^2\theta) = 0$ is used.
		4	

© Cambridge University Press & Assessment 2025

Question	Answer	Marks	Guidance
6(a)	State $f(x) \geqslant -3$ or $y \geqslant -3$ or clear equivalent	**B1**	Must be $\geqslant$ rather than >. Allow $\geqslant -3$ but not $x \geqslant -3$.
		1	
6(b)	Attempt to arrange to $x = ...$ in terms of y or equivalent	**M1**	Or in terms of x already. Sign errors only.
	Obtain $-3 + \sqrt{x+12}$	**A1**	Now in terms of x. A0 for $\pm$ in final answer. Condone $y = \ldots$
		2	
6(c)	Attempt expression for $gf(x)$	***M1**	Expect $2((x + 3)^2 - 12) - 5$. Sign errors only.
	Obtain $2(x+3)^2 - 29 = 69$	**A1**	OE E.g. $2x^2 + 12x - 80$ [= 0].
	Attempt solution of quadratic equation to find at least one value of x	**DM1**	No method needed.
	Obtain $x = 4$ only	**A1**	
	Alternative Method for Question 6(c)		
	Attempt solution of $g(x) = 69$ or evaluation of $g^{-1}(69)$	***M1**	Incorrect order of f^{-1} and then g^{-1} scores *M0.
	Obtain 37	**A1**	
	Attempt solution of $f(x) = \textit{their}\ 37$ or evaluation of $f^{-1}(\textit{their}\ 37)$	**DM1**	
	Obtain $x = 4$ only	**A1**	
		4	

© Cambridge University Press & Assessment 2025

Question	Answer	Marks	Guidance
7(a)	Obtain correct $\frac{1}{2}r^2 \times \frac{2}{3}\pi - \frac{1}{2}r^2 \sin\frac{2}{3}\pi$	**M1**	
	$0.614r^2$	**A1**	AG Greater accuracy is $0.61418...r^2$.
		2	
7(b)(i)	Obtain $\left[\frac{\mathrm{d}A}{\mathrm{d}r}\right] = 1.228\mathrm{r}$	**B1**	Or greater accuracy
	Use $\frac{\mathrm{d}A}{\mathrm{d}t} = \frac{\mathrm{d}A}{\mathrm{d}r} \times \frac{\mathrm{d}r}{\mathrm{d}t}$, or equivalent, with $r = 20$	**M1**	*'their'* $\frac{\mathrm{d}A}{\mathrm{d}r} \times 0.4$ with $r = 20$
	Obtain 9.8	**A1**	AWRT
		3	
7(b)(ii)	State or imply $\frac{2}{3}\pi r$ for length of arc AB	**B1**	Could be implied by $\frac{\mathrm{d}l}{\mathrm{d}\theta} = \frac{2\pi}{3}$.
	Differentiate and apply correct use of chain rule	**M1**	
	Obtain 0.84	**A1**	AWRT
		3	

© Cambridge University Press & Assessment 2025

Question	Answer	Marks	Guidance
8(a)	Integrate to obtain expression $\frac{1}{2}x^{\frac{3}{2}} \div \frac{3}{2}$	**B1**	OE
	Evaluate integral of the form $kx^{\frac{3}{2}}$ between limits 0 and 9 and subtract result from $\frac{27}{2}$	**M1**	
	Obtain $\frac{1}{3}x^{\frac{3}{2}}$ giving area under curve is 9 and shaded area is $\frac{9}{2}$	**A1**	
	Alternative Method for Question 8(a)		
	Obtain $x = 4y^2$	**B1**	
	Integrate to obtain expression of form $k\,y^3$ and evaluate between limits 0 and $\frac{3}{2}$	**M1**	
	Obtain $\frac{4}{3}y^3$ and hence shaded area is $\frac{9}{2}$	**A1**	
		3	
8(b)	Attempt to express x^2 in terms of y	**M1**	Condone ky^4. Attempting $\int y^2\,dx$ scores 0/4.
	Obtain or imply volume is $[\pi]\int 16y^4\,[dy]$	**A1**	
	Integrate to obtain ky^5 and evaluate between limits 0 and $\frac{3}{2}$	**M1**	
	Obtain $\frac{16}{5}y^5$ and hence $\frac{243}{10}\pi$ or exact equivalent	**A1**	Condone omission of π except for last mark.
		4	

© Cambridge University Press & Assessment 2025

Question	Answer	Marks	Guidance
9(a)	Attempt to express each of $S_2 - 1$, S_4, S_9 in terms of d (with at least one correct)	***M1**	Allow unsimplified.
	Obtain $S_2 - 1 = 3 + d$, $S_4 = 8 + 6d$, $S_9 = 18 + 36d$	**A1**	Must be simplified but can be implied by later work.
	Attempt equation in d by linking the three values	**DM1**	
	Obtain correct $(3 + d) + (18 + 36d) = 2(8 + 6d)$ or equivalent, and hence $d = -\frac{1}{5}$	**A1**	
		4	
9(b)	2 + 14 × *their d*	**M1**	
	Second progression: obtain either first term = $\frac{14}{5}$ or common difference = 4	**B1**	Must be from correct d in part **(a)**.
	Second progression: attempt to find 15th term	**M1**	Using a = 3 + *their d* from part **(a)**.
	Obtain 58.8, and hence difference = 59.6	**A1**	
		4	

© Cambridge University Press & Assessment 2025

Question	Answer	Marks	Guidance
10(a)	Attempt to substitute for x or y to produce quadratic equation in one variable	***M1**	Allow unsimplified.
	Obtain $5x^2 - 40x + 75 = 0$ or $5y^2 - 20 = 0$	**A1**	Or similarly simplified equivalent.
	Find two values of one variable	**B1**	
	Obtain coordinates $(3, 2)$ and $(5, -2)$	**B1**	
		4	
10(b)	Obtain centre $(0, -2)$ [and radius 5]	**B1**	
	Attempt area of ABC using mid-point of AB or $\frac{1}{2} \times \text{base} \times \text{height}$	**M1**	OE Use of correct formula for area of a triangle.
	Obtain $\frac{1}{2}\sqrt{20} \times \sqrt{20}$ or $\frac{1}{2} \times 5 \times 4$ (or equivalent) and hence 10	**A1**	
		3	

© Cambridge University Press & Assessment 2025

Question	Answer	Marks	Guidance
11(a)	Substitute 4 to obtain gradient of curve is $\frac{1}{10}$	**B1**	
	Attempt equation of normal using $(4, 3)$ and $-\frac{1}{m}$ for their gradient	**M1**	
	$y = -10x + 43$	**A1**	
		3	
11(b)	$\{-16x^{-3}\}\{+40(2x-3)^{-3}\}$	**B1 B1**	OE
	Substitute 4 to obtain $\frac{7}{100}$	**B1**	
		3	
11(c)	$[y =]\{-8x^{-1}\}\{+5(2x-3)^{-1}\}[+c]$	**B1 B1**	
	Substitute $x = 4$, $y = 3$ in an integrated expression to find value of c	**M1**	
	Obtain $3 = -2 + 1 + c$ and hence $y = -8x^{-1} + 5(2x-3)^{-1} + 4$	**A1**	OE For finding $c = 4$.
	Substitute $x = -1$ to obtain $q = 11$	**A1**	Not $y = 11$.
		5	

© Cambridge University Press & Assessment 2025