CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge International General Certificate of Secondary Education

MARK SCHEME for the March 2016 series

0606 ADDITIONAL MATHEMATICS

0606/12 Paper 12, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the March 2016 series for most Cambridge IGCSE® and Cambridge International A and AS Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.


CAMBRIDGE
International Examinations

Abbreviations

awrt answers which round to
cao correct answer only
dep dependent
FT follow through after error
isw ignore subsequent working
oe or equivalent
rot rounded or truncated
SC Special Case
soi seen or implied
www without wrong working

Question	Answer	Marks	Guidance
1	$ax+9=-2x^2+3x+1$ $2x^2+(a-3)x+8=0$	**M1**	for attempt to equate the line and the curve and obtain a 3 term quadratic equation
	For 2 distinct roots, $(a-3)^2>64$	**M1**	for use of the discriminant
	Critical values -5 and 11	**A1**	for critical values
	$a>11,\quad a<-5$	**A1**	for correct range
2	$a=-\frac{13}{6},\ b=0,\ c=1$	**B3**	**B1** for each
3	$\log_5\sqrt{x}+\log_{25}x=3$ $\frac{1}{2}\log_5 x+\frac{\log_5 x}{\log_5 25}=3$	**B1,B1**	**B1** for $\frac{1}{2}\log_5 x$ **B1** for $\frac{\log_5 x}{\log_5 25}$
	$\log_5 x=3$ $x=125$ cao	**B1**	for final answer
	Alternative scheme: $\frac{\log_{25}\sqrt{x}}{\log_{25}5}+\log_{25}x=3$	**B1**	for change of base
	$\frac{\frac{1}{2}\log_{25}x}{\log_{25}5}+\log_{25}x=3$	**B1**	for $\frac{1}{2}\log_{25}x$ (must be from correct work)
	$\log_{25}x=\frac{3}{2}$ $x=125$ cao	**B1**	for final answer

© Cambridge International Examinations 2016

Question	Answer	Marks	Guidance
4 (i)		**B1** **B1** **B1** **B1**	for a line in correct position for $(0, 2)$, $(2, 0)$ for correct shape for $y = \lvert 3 + 2x \rvert$, touching the x-axis for $(-1.5, 0)$, $(0, 3)$
(ii)	$2 - x = 3 + 2x$ leading to $x = -\frac{1}{3}$	**B1**	for $x = -\frac{1}{3}$
	$2 - x = -3 - 2x$ leading to $x = -5$	**M1** **A1**	for correct attempt to deal with 'negative' branch. for $x = -5$
	Alternative: $(2 - x)^2 = (3 + 4x)^2$ leading to $15x^2 + 28x + 5 = 0$ $x = -\frac{1}{3}$, $x = -5$	**M1** **A1,A1**	for equating and squaring to obtain a 3 term quadratic equation **A1** for each.
5 (a) (i)	${}^9P_6 = 60480$	**B1**	Must be evaluated
(ii)	${}^4P_2 \times {}^3P_2 \times 2 = 144$	**M1,A1**	**M1** for attempt a product of 3 perms
(iii)	840×2 1680	**B1,B1**	**B1** for either 840, or realising that there are 2 possible positions for the symbols
(b) (i)	${}^{10}C_6 \times {}^5C_3$ 2100	**M1** **A1**	for unsimplified form
(ii)	${}^8C_4 \times {}^4C_2$ 420	**M1** **A1**	for unsimplified form
6 (i)	$f(x) > 6$	**B1**	Allow **B1** for $y > 6$
(ii)	$f^{-1}(x) = \frac{1}{4}\ln(x - 6)$	**M1** **A1**	for a complete method must be $f^{-1}(x) =$ or $y = \ldots$
	Domain: $x > 6$ Range: $f^{-1}(x) \in \mathbb{R}$	**B1** **B1**	must be using the correct variable in both
(iii)	$f'(x) = 4e^{4x}$	**B1**	
(iv)	$6 + e^{4x} = 4e^{4x}$ leading to $x = \frac{1}{4}\ln 2$	**M1** **A1**	for a complete, correct method

© Cambridge International Examinations 2016

Question	Answer	Marks	Guidance
7 (i)	$f\left(\frac{1}{2}\right)=\frac{a}{8}+\frac{7}{4}-\frac{9}{2}+b \ \ (=0)$ $a+8b=22$	**M1**	for attempt at $f\left(\frac{1}{2}\right)$
	$8a+28-18+b=5(-a+7+9+b)$ $13a-4b=70$	**M1** **DM1**	for attempt at $f(2)=5f(-1)$ Allow if the 'wrong way' round for attempt to solve simultaneous equations
	leading to $a=6,\ b=2$	**A1**	**A1** for both
(ii)	$(2x-1)(3x^2+5x-2)$	**B2,1,0**	–1 each error
(iii)	$(2x-1)(3x-1)(x+2)$	**M1** **A1FT**	for attempt to factorise their quadratic factor must be 3 linear factors
8 (i)	$\lg y=\lg A+b\lg x$ Gradient = 1.2 so $b=1.2$	**B1** **M1** **A1**	may be implied by later work for attempt at gradient for $b=1.2$
	Intercept $=1.44$ $A=27.5$	**M1** **A1**	for attempt to find y-intercept for , allow awrt 28
(ii)	when $x=100$, $\lg x=2$ $\lg y=3.84$ (allow 3.8 to 3.9)	**M1** **A1**	for correct use of graph or equation
(iii)	when $y=8000,\ \lg 8000=3.9,\ \lg x=2.05$ leading to $x=113,\ 10^{2.05}$ or 112	**M1** **A1**	for correct use of graph or equation

© Cambridge International Examinations 2016

Question	Answer	Marks	Guidance
9 **(i)**	$\frac{7}{2}r^2\theta = \frac{1}{2}r^2(2\pi - \theta)$	**M1**	for a valid method
	$\theta = \frac{\pi}{4}$ oe	**A1**	allow in degrees
(ii)	$r + r + \frac{\pi}{4}r = 20$, leading to	**M1**	for valid method
	$r = 7.180(3..)$	**A1**	Must show enough accuracy to get **A1**
(iii)	Perimeter $= \frac{\pi}{4}r + 2r\tan\frac{\pi}{8}$ $= 5.6394 + 5.9484$ $= 11.6$	**B1,B1** **B1**	**B1** for arc length, **B1** for twice AC for 11.6
(iv)	Area $= (r \times AC) - \frac{1}{2}r^2\frac{\pi}{4}$ $= 21.356 - 20.246$ or equivalent method using triangles	**B1,B1**	**B1** for area of quadrilateral, allow unsimplified, **B1** for sector area
	$1.08 \leqslant$ Area $\leqslant 1.11$	**B1**	for area in given range
10 **(i)**	$x \times \frac{3}{2} \times 2(2x-1)^{\frac{1}{2}} + (2x-1)^{\frac{3}{2}}$	**B1** **M1** **A1**	for $\frac{3}{2} \times 2(2x-1)^{\frac{1}{2}}$ for attempt at differentiation of a product for all else correct
(ii)	$3\int x(2x-1)^{\frac{1}{2}}\,dx = x(2x-1)^{\frac{3}{2}} - \int(2x-1)^{\frac{3}{2}}\,dx$ $= x(2x-1)^{\frac{3}{2}} - \frac{1}{2} \times \frac{2}{5}(2x-1)^{\frac{5}{2}}$	**M1** **B1,B1**	for attempt to use part (i) **B1** for $x(2x-1)^{\frac{3}{2}}$, allow if divided by 3 **B1** for $\frac{1}{2} \times \frac{2}{5}(2x-1)^{\frac{5}{2}}$, allow if divided by 3
	$\int x(2x-1)^{\frac{1}{2}}\,dx = \frac{1}{3}(2x-1)^{\frac{3}{2}}\left(x - \frac{1}{5}(2x-1)\right)$ $= \frac{(2x-1)^{\frac{3}{2}}}{15}(3x+1)$	**M1** **DM1** **A1**	for taking out a common factor of $(2x-1)^{\frac{3}{2}}$ for attempt to obtain a linear factor
(iii)	$\left(\frac{1}{15} \times 4\right) - 0$	**M1** **A1FT**	for attempt to use limits correctly **FT** on *their* $\frac{px+q}{15}$

© Cambridge International Examinations 2016

Question	Answer	Marks	Guidance
11 (i)	$\dfrac{1}{\text{cosec}\,\theta-1}-\dfrac{1}{\text{cosec}\,\theta+1}=\dfrac{\text{cosec}\,\theta+1-\text{cosec}\,\theta+1}{\text{cosec}^2\theta-1}$	**M1**	for attempt to obtain a single fraction
		A1	all correct as shown
	$=\dfrac{2}{\cot^2\theta}$	**M1**	for use of correct identity
	$=2\tan^2\theta$	**A1**	for 'finishing off'
	Alternative scheme:		
	$\dfrac{1}{\text{cosec}\,\theta-1}-\dfrac{1}{\text{cosec}\,\theta+1}=\dfrac{\sin\theta}{1-\sin\theta}-\dfrac{\sin\theta}{1+\cos\theta}$	**M1**	for attempt to obtain a single fraction in terms of $\sin\theta$ only
	$=\dfrac{\left(\sin\theta+\sin^2\theta\right)-\left(\sin\theta-\sin^2\theta\right)}{1-\sin^2\theta}$	**A1**	all correct as shown
	$=\dfrac{2\sin^2\theta}{\cos^2\theta}$	**M1**	for use of correct identity
	$=2\tan^2\theta$	**A1**	for 'finishing off'
(ii)	$2\tan^2\theta=6+\tan\theta$ $\left(2\tan\theta+3\right)\left(\tan\theta-2\right)=0$	**M1**	for attempt to use (i), to obtain a quadratic equation and valid attempt to solve
	$\tan\theta=-\dfrac{3}{2}$, $\tan\theta=2$	**DM1**	for attempt to solve trig equation
	$\theta=63.4^\circ,\ 123.7^\circ,\ 243.4^\circ,\ 303.7^\circ$	**A1,A1**	for each 'pair'

© Cambridge International Examinations 2016