CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge International General Certificate of Secondary Education

MARK SCHEME for the May/June 2015 series

0606 ADDITIONAL MATHEMATICS

0606/11 Paper 1, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.


Cambridge is publishing the mark schemes for the May/June 2015 series for most Cambridge IGCSE®, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

CAMBRIDGE
International Examinations

Abbreviations

awrt answers which round to
cao correct answer only
dep dependent
FT follow through after error
isw ignore subsequent working
oe or equivalent
rot rounded or truncated
SC Special Case
soi seen or implied
www without wrong working

1	**(i)**	180° or π radians or 3.14 radians (or better)	**B1**	
	(ii)	2	**B1**	
	(iii) **(a)**		**B1**	$y = \sin 2x$ all correct
	(b)		**B1**	for either ↑↓↑starting at their highest value and ending at their lowest value Or a curve with highest value at $y = 3$ and lowest value at $y = -1$
			B1	completely correct graph
	(iv)	3	**B1**	
2	**(i)**	$\tan\theta = \dfrac{(8+5\sqrt{2})(4-3\sqrt{2})}{(4+3\sqrt{2})(4-3\sqrt{2})}$ $= \dfrac{32-24\sqrt{2}+20\sqrt{2}-30}{16-18}$ $= 1+2\sqrt{2}$ cao	**M1** **A1**	attempt to obtain $\tan\theta$ and rationalise. Must be convinced that no calculators are being used

© Cambridge International Examinations 2015

(ii)	$\sec^2\theta = 1+\tan^2\theta$ $= 1+\left(-1+2\sqrt{2}\right)^2$ $= 1+1-4\sqrt{2}+8$ $= 10-4\sqrt{2}$ **Alternative solution:** $AC^2 = \left(4+3\sqrt{2}\right)^2+\left(8+5\sqrt{2}\right)^2$ $= 148+104\sqrt{2}$ $\sec^2\theta = \dfrac{148+104\sqrt{2}}{\left(4+3\sqrt{2}\right)^2}$ $= \dfrac{148+104\sqrt{2}}{\left(4+3\sqrt{2}\right)^2}\times\dfrac{34-24\sqrt{2}}{34-24\sqrt{2}}$ $= 10-4\sqrt{2}$	**M1** **DM1** **A1** **M1** **DM1** **A1**	attempt to use $\sec^2\theta = 1+\tan^2\theta$, with *their* answer to **(i)** attempt to simplify, must be convinced no calculators are being used. Need to expand $\left(-1+2\sqrt{2}\right)^2$ as 3 terms
3 **(i)**	$64+192x^2+240x^4+160x^6$	**B3,2,1,0**	–1 each error
(ii)	$\left(64+192x^2+240x^4\right)\left(1-\dfrac{6}{x^2}+\dfrac{9}{x^4}\right)$ Terms needed $64-(192\times 6)+(240\times 9)$ $= 1072$	**B1** **M1** **A1**	expansion of $\left(1-\dfrac{3}{x^2}\right)^2$ attempt to obtain 2 or 3 terms using *their* **(i)**

© Cambridge International Examinations 2015

4	**(a)**	$\mathbf{X}^2 = \begin{pmatrix} 4-4k & -8 \\ 2k & -4k \end{pmatrix}$	**B2,1,0**	–1 each incorrect element
	(b)	Use of $\mathbf{AA}^{-1} = \mathbf{I}$ $\begin{pmatrix} a & 1 \\ b & 5 \end{pmatrix}\begin{pmatrix} \frac{5}{6} & -\frac{1}{6} \\ -\frac{2}{3} & \frac{1}{3} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	**M1**	use of $\mathbf{AA}^{-1} = \mathbf{I}$ and an attempt to obtain at least one equation.
		Any 2 equations will give $a = 2$, $b = 4$	**A1,A1**	
		Alternative method 1: $\frac{1}{5a-b}\begin{pmatrix} 5 & -1 \\ b & a \end{pmatrix} = \begin{pmatrix} \frac{5}{6} & -\frac{1}{6} \\ -\frac{2}{3} & \frac{1}{3} \end{pmatrix}$	**M1**	correct attempt to obtain $\mathbf{A}^{-1}$ and comparison of at least one term.
		Compare any 2 terms to give $a = 2$, $b = 4$	**A1,A1**	
		Alternative method 2: Inverse of $\frac{1}{6}\begin{pmatrix} 5 & -1 \\ -4 & 2 \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 4 & 5 \end{pmatrix}$	**M1** **A1,A1**	reasoning and attempt at inverse
5		$3x-1 = x(3x-1)+x^2-4$ or $y = \left(\frac{y+1}{3}\right)y + \left(\frac{y+1}{3}\right)^2 - 4$		
		$4x^2-4x-3=0$ or $4y^2-4y-35=0$ $(2x-3)(2x+1)=0$ or $(2y-7)(2y+5)=0$	**M1** **DM1**	equate and attempt to obtain an equation in 1 variable forming a 3 term quadratic equation and attempt to solve
		leading to $x = \frac{3}{2}, x = -\frac{1}{2}$ and	**A1**	x values
		$y = \frac{7}{2}, y = -\frac{5}{2}$	**A1**	y values
		Midpoint $\left(\frac{1}{2}, \frac{1}{2}\right)$	**B1**	for midpoint, allow anywhere
		Perpendicular gradient = $-\frac{1}{3}$	**M1**	correct attempt to obtain the gradient of the perpendicular, using AB
		Perp bisector: $y - \frac{1}{2} = -\frac{1}{3}\left(x - \frac{1}{2}\right)$	**M1**	straight line equation through the midpoint; must be convinced it is a perpendicular gradient.
		$(3y + x - 2 = 0)$	**A1**	allow unsimplified

© Cambridge International Examinations 2015

6	**(i)**	$f\left(\frac{1}{2}\right)=\frac{a}{8}-\frac{15}{4}+\frac{b}{2}-2=0$ leading to $a+4b=46$ $f(1)=a-15+b-2=5$ leading to $a+b=22$	**M1** **A1**	correct use of either $f\left(\frac{1}{2}\right)$ or f(1) paired correctly both equations correct (allow unsimplified)
		giving $b=8$ (AG), $a=14$	**M1,A1**	M1 for solution of equations A1 for both a and b. **AG** for b.
	(ii)	$(2x-1)(7x^2-4x+2)$	**M1,A1**	M1 for valid attempt to obtain g(x), by either observation or by algebraic long division.
	(iii)	$7x^2-4x+2=0$ has no real solutions as $b^2<4ac$ $16<56$	**M1** **A1**	use of b^2-4ac correct conclusion; must be from a correct g(x) or 2g(x) www
7	**(i)**	$\frac{dy}{dx}=\frac{(x-1)\frac{8x}{(4x^2+2)}-\ln(4x^2+3)}{(x-1)^2}$	**M1** **B1** **A1**	differentiation of a quotient (or product) correct differentiation of $\ln(4x^2+3)$ all else correct
		When $x=0$, $y=-\ln 3$ oe	**B1**	for y value
		$\frac{dy}{dx}=-\ln 3$ so gradient of normal is $\frac{1}{\ln 3}$ (allow numerical equivalent)	**M1**	valid attempt to obtain gradient of the normal
		normal equation $y+\ln 3=\frac{1}{\ln 3}x$ or $y=0.910x-1.10$, or $y=\frac{10}{11}x-\frac{11}{10}$ cao (Allow $y=0.91x-1.1$)	**M1** **A1**	attempt at normal equation must be using a perpendicular
	(ii)	when $x=0$, $y=-\ln 3$ when $y=0$, $x=(\ln 3)^2$ Area = ±0.66 or ±0.67 or awrt these or $\frac{1}{2}(\ln 3)^3$	**M1** **A1**	valid attempt at area

© Cambridge International Examinations 2015

8	**(i)**	Range for f: $y \geq 3$ Range for g: $y \geq 9$	**B1** **B1**	
	(ii)	$x = -2 + \sqrt{y-5}$	**M1**	attempt to obtain the inverse function
		$g^{-1}(x) = -2 + \sqrt{x-5}$ Domain of g^{-1}: $x \geq 9$	**A1** **B1**	Must be correct form for domain
		Alternative method: $y^2 + 4y + 9 - x = 0$	**M1**	attempt to use quadratic formula and find inverse
		$y = \dfrac{-4 + \sqrt{16 - 4(9-x)}}{2}$	**A1**	must have + not ±
	(iii)	Need $g(3e^{2x})$	**M1**	correct order
		$(3e^{2x} + 2)^2 + 5 = 41$	**DM1**	correct attempt to solve the equation
		or $9e^{4x} + 12e^{2x} - 32 = 0$ $(3e^{2x} - 4)(3e^{2x} + 8) = 0$		
		leading to $3e^{2x} + 2 = \pm 6$ so $x = \dfrac{1}{2}\ln\dfrac{4}{3}$	**M1**	dealing with the exponential correctly in order to reach a solution for x
		or $e^{2x} = \dfrac{4}{3}$ so $x = \dfrac{1}{2}\ln\dfrac{4}{3}$	**A1**	Allow equivalent logarithmic forms
		Alternative method: Using $f(x) = g^{-1}(41)$, $g^{-1}(41) = 4$	**M1**	correct use of g^{-1}
		leading to $3e^{2x} = 4$, so $x = \dfrac{1}{2}\ln\dfrac{4}{3}$	**DM1**	dealing with $g^{-1}(41)$ to obtain an equation in terms of e^{2x}
			M1 **A1**	dealing with the exponential correctly in order to reach a solution for x Allow equivalent logarithmic forms
	(iv)	$g'(x) = 6e^{2x}$ $g'(\ln 4) = 96$	**B1** **B1**	B1 for each

© Cambridge International Examinations 2015

9 **(i)**	$\frac{dy}{dx} = 3x^2 - 10x + 3$		
	When $x = 0$, for curve $\frac{dy}{dx} = 3$,		
	gradient of line also 3 so line is a tangent.	**A1**	comparing both gradients
	Alternate method:		
	$3x + 10 = x^3 - 5x^2 + 3x + 10$	**M1**	attempt to deal with simultaneous equations
	leading to $x^2 = 0$, so tangent at $x = 0$	**A1**	obtaining $x = 0$
(ii)	When $\frac{dy}{dx} = 0$, $(3x-1)(x-3) = 0$	**M1**	equating gradient to zero and valid attempt to solve
	$x = \frac{1}{3}$, $x = 3$	**A1,A1**	A1 for each
(iii)	Area $= \frac{1}{2}(10+19)3 - \int_0^3 x^3 - 5x^2 + 3x + 10\,dx$	**B1**	area of the trapezium
	$= \frac{87}{2} - \left[\frac{x^4}{4} - \frac{5x^3}{3} + \frac{3x^2}{2} + 10x\right]_0^3$	**M1**	attempt to obtain the area enclosed by the curve and the coordinate axes, by integration
	$= \frac{87}{2} - \left(\frac{81}{4} - 45 + \frac{27}{2} + 30\right)$	**A1**	integration all correct
		DM1	correct application of limits (must be using *their* 3 from **(ii)** and 0)
	$= 24.7$ or 24.8	**A1**	
	Alternative method:		
	Area $= \int_0^3 (3x+10) - \left(x^3 - 5x^2 + 3x + 10\right)dx$	**B1**	correct use of 'Y–y'
	$= \int_0^3 -x^3 + 5x^2\,dx$	**M1**	attempt to integrate
		A1	integration all correct
	$= \left[-\frac{x^4}{4} + \frac{5x^3}{3}\right]_0^3 = \frac{99}{4}$	**DM1**	correct application of limits
		A1	
10 **(a)**	$\sin^2 x = \frac{1}{4}$		
	$\sin x = (\pm)\frac{1}{2}$	**M1**	using $\text{cosec}\, x = \frac{1}{\sin x}$ and obtaining $\sin x = \ldots$
	$x = 30°, 150°, 210°, 330°$	**A1,A1**	A1 for one correct pair, A1 for another correct pair with no extra solutions

© Cambridge International Examinations 2015

(b)	$\left(\sec^2 3y - 1\right) - 2\sec 3y - 2 = 0$	**M1**	use of the correct identity
	$\sec^2 3y - 2\sec 3y - 3 = 0$ $(\sec 3y + 1)(\sec 3y - 3) = 0$ leading to $\cos 3y = -1$, $\cos 3y = \frac{1}{3}$	**M1**	attempt to obtain a 3 term quadratic equation in sec 3y and attempt to solve dealing with sec and 3y correctly
	$3y = 180°, 540°$ $\quad 3y = 70.5°, 289.5°, 430.5°$ $y = 60°, 180°, 23.5°, 96.5°, 143.5°$	**A1,A1** **A1**	A1 for a correct pair, A1 for a second correct pair, A1 for correct 5th solution and no other within the range
	Alternative 1:		
	$\sec^2 3y - 2\sec 3y - 3 = 0$	**M1**	use of the correct identity
	leading to $3\cos^2 3y + 2\cos 3y - 1$ $(3\cos y - 1)(\cos y + 1) = 0$	**M1** **M1**	attempt to obtain a quadratic equation in cos 3y and attempt to solve dealing with 3y correctly **A** marks as above
	Alternative 2:		
	$\frac{\sin^2 y}{\cos^2 y} - \frac{2}{\cos y} - 2 = 0$ $\left(1 - \cos^2 x\right) - 2\cos x - 2\cos^2 x = 0$	**M1**	use of the correct identity, $\tan y = \frac{\sin y}{\cos y}$ and $\sec y = \frac{1}{\cos y}$, then as before
(c)	$z - \frac{\pi}{3} = \frac{\pi}{3}, \frac{4\pi}{3}$	**M1**	correct order of operations
	$z = \frac{2\pi}{3}, \frac{5\pi}{3}$ or 2.09 or 2.1, 5.24	**A1,A1**	A1 for a correct solution A1 for a second correct solution and no other within the range

© Cambridge International Examinations 2015