Cambridge International Examinations
Cambridge International General Certificate of Secondary Education

ADDITIONAL MATHEMATICS **0606/21**

Paper 2 **May/June 2016**

MARK SCHEME

Maximum Mark: 80

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.


Cambridge is publishing the mark schemes for the May/June 2016 series for most Cambridge IGCSE®, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

© UCLES 2016

Abbreviations


awrt answers which round to
cao correct answer only
dep dependent
FT follow through after error
isw ignore subsequent working
oe or equivalent
rot rounded or truncated
SC Special Case
soi seen or implied
www without wrong working

Question	Answer	Marks	Guidance
1	$x^2 - 2x - 15$	**M1**	expands and rearranges to form a 3 term quadratic
	critical values −3 and 5	**A1**	not from wrong working
	$x < -3 \quad x > 5$	**A1**	mark final inequality; **A0** if spurious attempt to combine e.g. $5 < x < -3$
2 (a)		**B1**	It must be clear how the sets are nested
(b) (i)	$h \in P$	**B1**	Allow $\{m,a,t,h,s\}$ for P
(ii)	$n(P \cap Q) = 2$ cao	**B1**	
(iii)	{ t, h, s}	**B1**	
3 (i)	−2	**B1**	
(ii)	$-n$	**B1**	
(iii)	$\frac{\lg 5}{\log_5 10} = [(\lg y)^2]$ or $\frac{\lg 20 - \lg 4}{1 \big/ \lg 5} = [(\lg y)^2]$	**M1**	One log law used correctly
	correct completion to $(\lg 5)^2$ isw	**A1**	answer only does not score
(iv)	$[\log_r]6x^2 = [\log_r]600$	**B1**	Condone base missing
	$x = 10$ only	**B1**	


© Cambridge International Examinations 2016

Question	Answer	Marks	Guidance
4 (i)	$\frac{\pi}{3}$ isw	B1	
(ii)	[Area triangle ABC =] $\frac{1}{2}\times 10^2 \times \sin\left(their\ \frac{\pi}{3}\right)$ oe	M1	seen or implied by $25\sqrt{3}$ or 43.3(0...)
	[Area 1 sector =] $\frac{1}{2}\times 5^2 \times their\ \frac{\pi}{3}$ oe or $\pi \times 5^2 \times \frac{their 60°}{360}$	M1	seen or implied by $\frac{25\pi}{6}$ or 13.0(8...) or 13.09
	Complete correct plan	M1	e.g. *their* triangle – 3(*their* sector)
	4.03(1...) or $25\sqrt{3} - \frac{25\pi}{2}$ isw	A1	Units not required
5 (a)	$\frac{\sqrt{8}}{\left(\sqrt{7}-\sqrt{5}\right)} \times \frac{\left(\sqrt{7}+\sqrt{5}\right)}{\left(\sqrt{7}+\sqrt{5}\right)}$ and attempt to multiply	M1	
	$\frac{\sqrt{56}+\sqrt{40}}{2}$ oe	A1	not from wrong working
	$\sqrt{14}+\sqrt{10}$	A1	
(b)	$q^2 + 4q\sqrt{3} + 12$ soi	B1	
	$28 = q^2 + 12$ oe	M1	can be implied by 4 and 16 or –4 and –16
	$q =$ 4, –4 $p =$ 16, –16	A1	all values
6 (i)	$4(x+1)^2 - 9$	B3,2,1,0	one mark for each of p, q, r correct in a correctly formatted expression; allow correct equivalent values; If **B0** then **SC2** for $4(x+1)-9$ or **SC1** for correct 3 values seen in incorrect format e.g. $4(x+1x)-9$ or $4(x^2+1)-9$ or for a correct completed square form of the original expression in a different but correct format. e.g. $2(\sqrt{2}x+\sqrt{2})^2-9$

© Cambridge International Examinations 2016

Question	Answer	Marks	Guidance
(ii)	(–1, 9)	B2FT	B1FT $(-q\ ,-r)$ $r<0$ for each correct coordinate
(iii)			
		B1	Correct symmetric W shape with cusps on x-axis
		B1	y-intercept marked at 5 only or coords indicated on graph
		B1	x-intercepts marked at –2.5 and 0.5 only x-axis or coords indicated on graph or close by
7 (i) (a)	$\mathbf{q}-\mathbf{p}$	B1	
(b)	$2\mathbf{q}-2\mathbf{p}$ or $2(\mathbf{q}-\mathbf{p})$	B1	
(ii)	The points are collinear oe	B1	
	$\overrightarrow{PQ}$ is a (scalar) multiple of $\overrightarrow{QR}$ **and** they have a point in common. oe	B1	Condone $\overrightarrow{PQ}$ is parallel to $\overrightarrow{QR}$ **and** …
(iii)	[$\overrightarrow{OR}$ =] $4\mathbf{i}-3\mathbf{j}$ oe soi	B1	
	$\sqrt{4^2+(-3)^2}$ (=5)	M1	condone $\sqrt{4^2+3^2}$; may be implied by correct answer or correct FT answer
	$\frac{1}{5}(4\mathbf{i}-3\mathbf{j})$ oe	A1	
8 (a) (i)	$a^4+4a^3b+6a^2b^2+4ab^3+b^4$ final answer	B2,1,0	–1 each error/omission
(ii)	$6(2x)^2\left(\frac{1}{5x}\right)^2$ soi	M1	Could be in full expansion
	$\frac{24}{25}$ or 0.96 isw	A1	Must be explicitly identified
(b)	$\frac{1}{8}\left(\frac{n(n-1)(n-2)}{6}\right)=\frac{5n}{12}$ soi leading to a cubic or quadratic ($n^2-3n-18=0$)	M1	Must attempt to expand and remove fractions
	Solves *their* quadratic [$(n-6)(n+3)$]	M1	must have come from a valid attempt
	[n =] 6 only, not from wrong working	A1	Must be n if labelled

© Cambridge International Examinations 2016

Question	Answer	Marks	Guidance
9 (a)	$a = 2 \quad b = 4 \quad c = -2$	B3	**B1** for each correct value
(b) (i)		B3,2,1,0	sinusoidal curve symmetrical about y-axis clear intent to have amplitude of 2 2 cycles If not fully correct max **B2**
(ii)	$-\frac{\pi}{2}, -\frac{\pi}{6}, \frac{\pi}{6}, \frac{\pi}{2}, -\frac{\pi}{3}, \frac{\pi}{3}$ cao	B2	**B1** for any 4 correct
10 (a) (i)	$2 \times 4!$ or $\frac{2}{5} \times 5!$ oe	M1	
	48	A1	
(ii)	${}^5P_3$ or $\frac{5!}{2!}$ or $5 \times 4 \times 3$ oe	M1	
	60	A1	
(b) (i)	$4 \times 2[!] \times 3$oe	M1	Correct first step implied by a correct product of two elements
	24	A1	
(ii)	3! or 3×3 seen	M1	
	18	A1	
11 (i)	$\frac{3x^2}{2} - \frac{2x^{5/2}}{5}(+c)$ isw	B1+B1	
(ii)	(9, 0) oe	B1	Not just $x = 9$
(iii)	Substitute (3, 9) into **both** lines Or solves simultaneously ($6x = 27 - 3x$ oe) to get $x = 3$, $y = 9$	B1	$3 \times 3 = 9$ **and** $\frac{27 - 3 \times 3}{2} = 9$

© Cambridge International Examinations 2016

Question	Answer	Marks	Guidance
(iv)	$\left[\text{Area } AOB = \right]\frac{1}{2}\times 9\times 9$ oe ($\frac{81}{2}$ or 40.5)	**M1**	Uses *their* (ii). May split into 2 triangles (13.5 and 27). May integrate. Must be a complete method.
	$their\left[\frac{3(9)^2}{2}-\frac{2(9)^{\frac{5}{2}}}{5}\right]-[0]$ (= 24.3)	**M1**	lower limit may be omitted but must be correct if seen
	$their\frac{81}{2}-their\frac{243}{10}$	**M1**	must be from genuine attempts at area of triangle and area under curve
	16.2	**A1**	
12 (i)	$\left[\frac{dy}{dx}=\right]\frac{2(x-1)-(2x-5)}{(x-1)^2}$	**M1A1**	Allow slips in $\frac{du}{dx}$ and $\frac{dv}{dx}$ but must be explicit. Allow $(x-1)^2=x^2-2x+1$
	– 12 isw	**B1**	
	ALT using $y=\frac{-12x^2+14x-5}{x-1}$		
	$-24x+14$	**B1**	
	$\left[\frac{dy}{dx}=\right]\frac{(x-1)(-24x+14)-(-12x^2+14x-5)}{(x-1)^2}$	**M1**	
		A1FT	**FT** on their derivative of 3 term quadratic
(ii)	$\left[\frac{d^2y}{dx^2}=\right]k(x-1)^{-3}$	**M1**	No additional terms
	$k = -6$ isw	**A1**	

© Cambridge International Examinations 2016

Question	Answer	Marks	Guidance
(iii)	*their* $\left[\frac{3}{(x-1)^2}-12\right]=0$ and find a value for x	**M1**	$12x^2-24x+9=0$ oe $(2x-3)(2x-1)=0$ oe
	$x = 0.5$ and $x = 1.5$	**A1**	
	$y = 2$ and $y = -22$	**A1**	if **A0 A0** then **A1** for a correct (x, y) pair
	$\frac{-6}{(-0.5)^3}>0$ therefore min when $x = 0.5$ oe	**B1**	or $\left[\frac{-6}{(-0.5)^3}=\right]48$ therefore min when $x = 0.5$ oe
	$\frac{-6}{(0.5)^3}<0$ therefore max when $x = 1.5$ oe	**B1**	or $\left[\frac{-6}{(0.5)^3}=\right]-48$ therefore max when $x = 1.5$ oe **M1A1** is possible from other methods

© Cambridge International Examinations 2016