CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge International General Certificate of Secondary Education

MARK SCHEME for the October/November 2014 series

0606 ADDITIONAL MATHEMATICS

0606/23 Paper 2, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2014 series for most Cambridge IGCSE®, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

CAMBRIDGE
International Examinations

1	**(i)**	$f(2)=0 \rightarrow 3(2)^3+8(2)^2-33(2)+p=0$ correct working to $p = 10$ AG method for quadratic factor $f(x)=(x-2)(3x^2+14x-5)$	**M1** **A1** **M1** **A1**	
	(ii)	$f(x)=(x-2)(3x-1)(x+5)$ $f(x)=0 \rightarrow x=2,\ -5,\ \frac{1}{3}$	**M1** **A1**	factorise or solve quadratic factor = 0
2	**(i)**	${}^{12}C_4=495$	**B1**	
	(ii)	${}^7C_2\times{}^5C_2=21\times10$ $=210$	**M1** **A1**	
	(iii)	not K and B = ${}^6C_2\times{}^4C_1=15\times4=60$ K and not B = ${}^6C_1\times{}^4C_2=6\times6=36$ 60 + 36 96 OR K and B = ${}^6C_1\times{}^4C_1=6\times4=24$ not K and not B = ${}^6C_2\times{}^4C_2=15\times6=90$ 210 – 90 – 24 96	**B1** **B1** **M1** **A1** **B1** **B1** **M1** **A1**	
3	**(i)**	C is (1, 6) D is $(1,6)+(12,9)$ = (13, 15)	**B1** **M1** **A1ft**	
	(ii)	gradient of $CD=\frac{15-6}{13-1}\left(=\frac{3}{4}\right)$ gradient of $AB=\frac{10-2}{-2-4}\left(=\frac{8}{-6}=\frac{-4}{3}\right)$ $\frac{3}{4}\times\frac{-4}{3}=-1$ lines are perpendicular	**B1ft** **B1** **B1**	 correct completion www
	(iii)	area $=\frac{1}{2}\times AB\times CD=\frac{1}{2}\times10\times15$ =75 or array method	**M1** **A1**	good attempt at two relevant lengths for $\frac{1}{2}$ base × height method

© Cambridge International Examinations 2014

4	**(i)**	$2000 = 1000e^{a+b} \rightarrow a+b = \ln 2$	**B1**	
	(ii)	$3297 = 1000e^{2a-b} \rightarrow 2a+b$ $= \ln 3.297$ oe	**M1** **A1**	substitution of 2, 3297 and rearrange
	(iii)	Solve for one value $a = 0.5$ and $b = 0.193$ or 0.19	**M1** **A1**	
	(iv)	$n = 10 \quad P = 1000e^{5.193}$ = \$180 000.	**M1** **A1**	
5	**(i)**	$\overrightarrow{OX} = \mu(a+b)$	**B1**	
	(ii)	$\overrightarrow{RP} = b-3a$ or $\overrightarrow{RX} = \lambda(b-3a)$ oe $\overrightarrow{OX} = 3a+\lambda(b-3a)$	**B1** **B1**	
	(iii)	$\overrightarrow{OX} = \overrightarrow{OX}$ and equate both coefficients $\mu = 3-3\lambda \quad \mu = \lambda$ $\mu = \lambda = 0.75$ $\frac{RX}{XP} = 3$ or 3:1	**M1** **A1** **A1ft**	$\frac{\lambda}{1-\lambda}$
6	**(i)**	$m = 4$ equation of line is $\frac{\ln y - 39}{3^x - 9} = \frac{39-19}{9-4}$ $\ln y = 4(3^x)+3$	**B1** **M1** **A1ft**	forms equation of line ft only on their gradient
	(ii)	$x = 0.5 \rightarrow \ln y = 4\sqrt{3}+3 = 9.928$ $y = 20\,500$	**M1** **A1**	correct expression for ln$y$
	(iii)	Substitutes y and rearrange for 3^x Solve $3^x = 1.150$ $x = 0.127$	**M1** **M1** **A1**	

© Cambridge International Examinations 2014

7	**(i)**	$x = \frac{2}{y} + 1 \quad \rightarrow \quad y = \frac{2}{x-1}$	**M1**	any valid method
		$f^{-1}(x) = \frac{2}{x-1}$	**A1**	
	(ii)	$gf(x) = \left(\frac{2}{x} + 1\right)^2 + 2$	**B2/1/0**	–1 each error
	(iii)	$fg(x) = \frac{2}{x^2+2} + 1$	**B2/1/0**	–1 each error
	(iv)	$ff(x) = \frac{2}{\frac{2}{x}+1} + 1 = \frac{2x}{x+2} + 1$	**M1**	correct starting expression
		$= \frac{3x+2}{x+2}$	**A1**	correct algebra to given answer
		$\frac{3x+2}{x+2} = x \quad \rightarrow \quad x^2 - x - 2 = 0$	**M1**	form and solve 3 term quadratic
		$(x-2)(x+1) = 0$ $x = 2$ only	**A1**	
8	**(i)**	$v = C + K\sin 2t \qquad C \neq 0$	**M1**	
		$v = 5 + 6\sin 2t$	**A1**	
		$a = 12\cos 2t$	**A1ft**	
	(ii)	$a = 0 \quad \rightarrow \quad \cos 2t = 0$ and solve	**M1**	set $a = 0$ and solve for t
		$t = \frac{\pi}{4}$ or 0.785 or 0.79	**A1**	
		$v = 5 + 6\sin\frac{\pi}{2} = 11$	**A1ft**	ft only on K
	(iii)	$v = 2 \quad \rightarrow \quad \sin 2t = -\frac{1}{2}$ and solve	**M1**	set $v = 2$ and solve for t
		$t = \frac{7\pi}{12}$ or 1.83 – 1.84	**A1**	
		$a = 12\cos\frac{7\pi}{6} = -6\sqrt{3}$ or –10.4	**A1**	

© Cambridge International Examinations 2014

9	**(i)**	$\frac{dy}{dx} = 4 - \frac{1}{(x-2)^2}$	**B1**	
		$\frac{dy}{dx} = 0 \rightarrow (x-2)^2 = \frac{1}{4}$ $(4x^2 - 16x + 15 = 0)$	**M1**	solve 3 term quadratic from $\frac{dy}{dx} = 0$
		$x = 2.5$ or 1.5	**A1**	x values or 1 pair
		$y = 12$ or 4	**A1**	y values or 1 pair
		$\frac{d^2y}{dx^2} = 2(x-2)^{-3}$	**M1**	use $\frac{d^2y}{dx^2}$ with solution from $\frac{dy}{dx} = 0$
		$x = 2.5 \rightarrow \frac{d^2y}{dx^2} > 0 \rightarrow$ minimum $x = 1.5 \rightarrow \frac{d^2y}{dx^2} < 0 \rightarrow$ maximum	**A1**	both identified www
	(ii)	$x = 3 \rightarrow \frac{dy}{dx} = 3$	**B1**	
		Use $m_1 m_2 = -1$ for gradient normal from gradient tangent	**M1**	must use numerical values
		Eqn of normal : $\frac{y-13}{x-3} = -\frac{1}{3}$	**A1ft**	
		Intersection of norm and curve		
		$14 - \frac{x}{3} = 4x + \frac{1}{x-2}$	**M1**	equation and attempt to simplify
		$13x^2 - 68x + 87 = 0$	**DM1**	attempt to solve 3 term quadratic
		$x = \frac{29}{13}$ or 2.23	**A1**	
10	**(i)**	$\text{LHS} = \frac{1+\cos x + 1 - \cos x}{(1-\cos x)(1+\cos x)}$	**B1**	correct fraction
		$= \frac{2}{1-\cos^2 x}$	**B1**	correct evaluation
		$= \frac{2}{\sin^2 x} = \text{RHS}$	**B1**	use of $1 - \cos^2 x = \sin^2 x$ and completion of fully correct proof
	(ii)	$2\text{cosec}^2 x = 8$	**M1**	identity used
		$\sin^2 x = \frac{1}{4}$	**A1**	
		$\sin x = \pm\frac{1}{2}$	**A1**	
		$x = 30^\circ, 150^\circ, 210^\circ, 330^\circ$	**A1**	

© Cambridge International Examinations 2014