CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge International General Certificate of Secondary Education

MARK SCHEME for the October/November 2015 series

0606 ADDITIONAL MATHEMATICS

0606/12 Paper 1, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2015 series for most Cambridge IGCSE®, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

CAMBRIDGE
International Examinations

Abbreviations

awrt answers which round to
cao correct answer only
dep dependent
FT follow through after error
isw ignore subsequent working
oe or equivalent
rot rounded or truncated
SC Special Case
soi seen or implied
www without wrong working

1	$kx^2 + (2k-8)x + k = 0$	M1	for attempt to obtain a 3 term quadratic in the form $ax^2 + bx + c = 0$,where b contains a term in k and a constant
	$b^2 - 4ac > 0$ so $(2k-8)^2 - 4k^2 (>0)$	DM1	for use of $b^2 - 4ac$
	$4k^2 - 32k + 64 - 4k^2 (>0)$	DM1	for attempt to simplify and solve for k
	leading to $k < 2$ only	A1	A1 must have correct sign
2	$\left(\frac{dy}{dx}\right) = -5x(+c)$	M1	for attempt to integrate, do not penalise omission of arbitrary constant.
	When $x = -1$, $\frac{dy}{dx} = 2$ leading to		
	$\frac{dy}{dx} = -5x - 3$	A1	Must have $\frac{dy}{dx} = ...$
	$y = -\frac{5x^2}{2} - 3x + d$	DM1	for attempt to integrate *their* $\frac{dy}{dx}$, but penalise omission of arbitrary constant.
	When $x = -1, y = 3$ leading to		
	$y = \frac{5}{2} - \frac{5x^2}{2} - 3x$	A1	
	Alternative scheme:		
	$y = ax^2 + bx + c$ so $\frac{dy}{dx} = 2ax + b$	M1	for use of $y = ax^2 + bx + c$, differentiation and use of conditions to give an equation in a and b
	When $x = -1$, $\frac{dy}{dx} = 2$		
	so $-2a + b = 2$	A1	for a correct equation
	$\frac{d^2y}{dx^2} = 2a$	DM1	for a second differentiation to obtain a
	so $a = -\frac{5}{2}$, $b = -3$, $c = \frac{5}{2}$	A1	for a, b and c all correct

© Cambridge International Examinations 2015

3	$\sqrt{(\sec^2\theta - 1)} + \sqrt{(\text{cosec}^2\theta - 1)} = \sec\theta\,\text{cosec}\,\theta$		
	LHS $= \tan\theta + \cot\theta$	B1	may be implied by the next line
	$= \dfrac{\sin\theta}{\cos\theta} + \dfrac{\cos\theta}{\sin\theta}$	B1	for dealing with $\tan\theta$ and $\cot\theta$ in terms of $\sin\theta$ and $\cos\theta$
	$= \dfrac{\sin^2\theta + \cos^2\theta}{\sin\theta\cos\theta}$	M1	for attempt to obtain as a single fraction
	$= \dfrac{1}{\sin\theta\cos\theta}$	M1	for the use of $\sin^2\theta + \cos^2\theta = 1$ in correct context
	$= \sec\theta\,\text{cosec}\,\theta$	A1	Must be convinced as AG
	Alternate scheme:		
	LHS $= \tan\theta + \cot\theta$		
	$= \tan\theta + \dfrac{1}{\tan\theta}$	B1	may be implied by subsequent work
	$= \dfrac{\tan^2\theta + 1}{\tan\theta}$	M1	for attempt to obtain as a single fraction
	$= \dfrac{\sec^2\theta}{\tan\theta}$	B1	for use of the correct identity
	$= \dfrac{\sec\theta}{\tan\theta} \times \sec\theta$	M1	for 'splitting' $\sec^2\theta$
	$= \text{cosec}\,\theta\sec\theta$	A1	Must be convinced as AG
4 **(a)** **(i)**	28	B1	
(ii)	20160	B1	
(iii)	$6 \times (5 \times 4 \times 3)$ oe to give 360 $6 \times (5 \times 4 \times 3) \times 2$	B1	for realising that the music books can be arranged amongst themselves and consideration of the other 5 books
	$= 720$	B1	for the realisation that the above arrangement can be either side of the clock.
(b)	Either $^{10}C_6 - {}^7C_6 = 210 - 7$	B1, B1	B1 for $^{10}C_6$, B1 for $^7C_6$
	$= 203$	B1	
	Or 1W 5M = 63 2W 4M = 105	B1	for 1 case correct, must be considering more than 1 different case, allow *C* notation
	3W 3M = 35	B1	for the other 2 cases, allow *C* notation
	Total = 203	B1	for final result

© Cambridge International Examinations 2015

5	**(i)**	$\frac{dy}{dx} = (x-3)\frac{4x}{2x^2+1} + \ln(2x^2+1)$ when $x = 2$, $\frac{dy}{dx} = -\frac{8}{9} + \ln 9$ oe or 1.31 or better	B1 M1 A1 A1	for correct differentiation of ln function for attempt to differentiate a product for correct product, terms must be bracketed where appropriate for correct final answer
	(ii)	$\partial y \approx$ (answer to **(i)**) × 0.03 $= 0.0393$, allow awrt 0.039	M1 A1FT	for attempt to use small changes follow through on *their* numerical answer to **(i)** allow to 2 sf or better
6	**(i)**	$A \cap B = \{3\}$	B1	
	(ii)	$A \cup C = \{1, 3, 5, 6, 7, 9, 11, 12\}$	B1	
	(iii)	$A' \cap C = \{1, 5, 7, 11\}$	B1	
	(iv)	$(D \cup B)' = \{1, 9\}$	B1	
	(v)	Any set containing up to 5 positive even numbers $\leqslant 12$	B1	
7	**(i)**	Gradient $= \frac{0.2}{0.8} = 0.25$ $b = 0.25$	M1 A1	for attempt to find the gradient
		Either $6 = 0.25(2.2) + c$ Or $5.8 = 0.25(1.4) + c$ leading to $A = 233$ or $e^{5.45}$	M1 A1	for a correct substitution of values from either point and attempt to obtain c or solution by simultaneous equations dealing with $c = \ln A$
		Alternative schemes: Either: $6 = b(2.2) + c$; $5.8 = b(1.4) + c$ Or: $e^6 = A\left(e^{2.2}\right)^b$; $e^{5.8} = A\left(e^{1.4}\right)^b$	M1	for 2 simultaneous equations as shown
			DM1	for attempt to solve to get at least one solution for one unknown
		Leading to $A = 233$ or $e^{5.45}$ and $b = 0.25$	A1, A1	A1 for each
	(ii)	Either $y = 233 \times 5^{0.25}$ Or $\ln y = 0.25 \ln 5 + \ln 233$ leading to $y = 348$	M1 A1	for correct use of either equation in attempt to obtain y using *their* value of A and of b found in **(i)**

© Cambridge International Examinations 2015

8	$\dfrac{dy}{dx} = \dfrac{2(x^2+5)^{\frac{1}{2}} - \frac{1}{2}(2x)(x^2+5)^{-\frac{1}{2}}(2x-1)}{x^2+5}$ or $\dfrac{dy}{dx} = 2(x^2+5)^{-\frac{1}{2}} - \frac{1}{2}(2x)(x^2+5)^{-\frac{3}{2}}(2x-1)$	B1 M1 A1	for $\frac{1}{2}(2x)(x^2+5)^{-\frac{1}{2}}$ for a quotient or $-\frac{1}{2}(2x)(x^2+5)^{-\frac{3}{2}}$ for a product allow if either seen in separate working for attempt to differentiate a quotient or a **correct** product for all correct, allow unsimplified
	When $x=2, y=1$ and $\dfrac{dy}{dx} = \dfrac{4}{9}$ (allow 0.444 or 0.44)	B1, B1	B1 for each
	Equation of tangent: $y-1=\dfrac{4}{9}(x-2)$ $(9y = 4x+1)$	M1 A1	for attempt at straight line, must be tangent using *their* gradient and y allow unsimplified.
9 (i)	$\dfrac{2}{3}(4+x)^{\frac{3}{2}}$ $(+c)$	B1,B1	B1 for $k(4+x)^{\frac{3}{2}}$ only, B1 for $\dfrac{2}{3}(4+x)^{\frac{3}{2}}$ only Condone omission of c
(ii)	Area of trapezium $= \left(\dfrac{1}{2}\times 5\times 5\right)$ $= 12.5$	M1 A1	for attempt to find the area of the trapezium
	Area $= \left[\dfrac{2}{3}(4+x)^{\frac{3}{2}}\right]_0^5 - \left(\dfrac{1}{2}\times 5\times 5\right)$	M1	for correct use of limits using $k(4+x)^{\frac{3}{2}}$ only (must be using 5 and 0)
	$= \left(\dfrac{2}{3}\times 27\right) - \dfrac{16}{3} - \dfrac{25}{2}$	A1	for $18 - \dfrac{16}{3}$ or equivalent
	$= \dfrac{1}{6}$ or awrt 0.17	A1	
	Alternative scheme:		
	Equation of AB $y = \dfrac{1}{5}x + 2$	M1	for a correct attempt to find the equation of AB
	Area $= \int_0^5 \sqrt{4+x} - \left(\dfrac{1}{5}x+2\right)dx$	M1	for correct use of limits using $k(4+x)^{\frac{3}{2}}$ only (must be using 5 and 0)
	$= \left[\dfrac{2}{3}(4+x)^{\frac{3}{2}} - \dfrac{x^2}{10} - 2x\right]_0^5$		
	$= \left(\dfrac{2}{3}\times 27\right) - \dfrac{16}{3} - \dfrac{25}{2}$	A1	for $18 - \dfrac{16}{3}$ or equivalent
	$= \dfrac{1}{6}$ or awrt 0.17	A1 A1	for 12.5 or equivalent

© Cambridge International Examinations 2015

10 (i)	All sides are equal to the radii of the circles which are also equal	B1	for a convincing argument
(ii)	Angle $CBE = \frac{2\pi}{3}$	B1	must be in terms of π, allow 0.667π, or better
(iii)	$DE = 10\sqrt{3}$	M1	for correct attempt to find *DE* using *their* angle *CBE*
		A1	for correct *DE*, allow 17.3 or better
	Arc $CE = 10 \times \frac{2\pi}{3}$	M1	for attempt to find arc length with *their* angle *CBE* (20.94)
	Perimeter $= 20 + 10\sqrt{3} + \frac{20\pi}{3}$	M1	for 10 + 10 + *DE* + an arc length
	$= 58.3$ or 58.2	A1	allow unsimplified
(iv)	Area of sector: $\frac{1}{2} \times 10^2 \times \frac{2\pi}{3} = \frac{100\pi}{3}$	M1	for sector area using *their* angle *CBE* allow unsimplified, may be implied
	Area of triangle: $\frac{1}{2} \times 10^2 \times \sin\frac{2\pi}{3} = 25\sqrt{3}$	M1	for triangle area using *their* angle *DBE* which must be the same as *their* angle *CBE*, allow unsimplified, may be implied
	Area $= \frac{100\pi}{3} + 25\sqrt{3}$ or awrt 148	A1	allow in either form

© Cambridge International Examinations 2015

11	**(a)** **(i)**	$(x+3)^2 - 5$	B1, B1	B1 for 3, B1 for -5
	(ii)	$y \geqslant 4$ or $f \geqslant 4$	B1	Correct notation or statement must be used
	(iii)	$y = \sqrt{x+5} - 3$	M1	for a correct attempt to find the inverse function
			A1	must be in the correct form and positive root only
		Domain $x \geqslant 4$	B1FT	Follow through on *their* answer to **(ii),** must be using x
	(b)	$h^2g(x) = h^2(e^x)$	M1	for correct order
		$= h(5e^x + 2)$	M1	for dealing with h^2
		$= 25e^x + 12$		
		$25e^x + 12 = 37,$	DM1	for solution of equation (dependent on both previous M marks)
		leading to $x = 0$	A1	
		Alternative scheme 1:		
		$hg(x) = h^{-1}(37)$	M1	for correct order
		$h^{-1}(37) = 7$	M1	for dealing with $h^{-1}(37)$
		$5e^x + 2 = 7,$	DM1	for solution of equation (dependent on both previous M marks)
		leading to $x = 0$	A1	
		Alternative scheme 2:		
		$g(x) = h^{-2}(37)$	M1	for correct order
		$h^{-2}(37) = 1$	M1	for dealing with $h^{-2}(37)$
		$e^x = 1,$	DM1	for solution of equation (dependent on both previous M marks)
		leading to $x = 0$	A1	

© Cambridge International Examinations 2015

12	$x^2 + 6x - 16 = 0$ or $y^2 + 10y - 75 = 0$ leading to $(x+8)(x-2) = 0$ or $(y-5)(y+15) = 0$	M1 DM1	for attempt to obtain a 3 term quadratic in terms of one variable only for attempt to solve quadratic equation
	so $x = 2, y = 5$ and $x = -8, y = -15$	A1, A1	A1 for each 'pair' of values.
	Midpoint $(-3, -5)$	B1	
	Gradient $= 2$, so perpendicular gradient $= -\frac{1}{2}$ Perpendicular bisector: $y + 5 = -\frac{1}{2}(x+3)$ $(2y + x + 13 = 0)$	M1 M1	for attempt at straight line equation, must be using midpoint and perpendicular gradient for use of $y = 0$ in *their* line equation (but not $2x - y + 1 = 0$)
	Point C $(-13, 0)$		
	Area $= \frac{1}{2}\begin{vmatrix} -13 & 2 & -8 & -13 \\ 0 & 5 & -15 & 0 \end{vmatrix}$ $= 125$	M1 A1	for correct attempt to find area, may be using *their* values for A, B and C (C must lie on the x-axis)
	Alternative method for area: $CM^2 = 125,\ AB^2 = 500$ Area $= \frac{1}{2} \times \sqrt{125} \times \sqrt{500}$ $= 125$	M1 A1	for correct attempt to find area may be using *their* values for A, B and C

© Cambridge International Examinations 2015