CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge International General Certificate of Secondary Education

MARK SCHEME for the October/November 2015 series

0606 ADDITIONAL MATHEMATICS

0606/23 Paper 2 , maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2015 series for most Cambridge IGCSE®, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

CAMBRIDGE
International Examinations

Abbreviations

awrt answers which round to
cao correct answer only
dep dependent
FT follow through after error
isw ignore subsequent working
oe or equivalent
rot rounded or truncated
SC Special Case
soi seen or implied
www without wrong working

1	$y = x^3 + 3x^2 - 5x - 7$ $\frac{dy}{dx} = 3x^2 + 6x - 5$ $x = 2 \rightarrow \frac{dy}{dx} = 19$ $y = 3$ eqn of tangent: $\frac{y-3}{x-2} = 19 \quad \rightarrow \quad (y = 19x - 35)$	M1 A1 A1FT B1 A1FT	Differentiate on *their* $\frac{dy}{dx}$
2	$2x + k + 2 = 2x^2 + (k+2)x + 8$ $2x^2 + kx + 6 - k \quad (= 0)$ $b^2 - 4ac = k^2 - 4 \times 2(6-k)$ $k^2 + 8k - 48 \quad (> 0)$ $(k+12)(k-4) \quad (> 0)$ $k < -12$ or $k > 4$	M1 A1 M1 DM1 A1 A1	eliminate y or x correct quadratic use discriminant attempt to solve 3 term quadratic $k = -12$ and $k = 4$
3 (a)	$\frac{dy}{dx} = \frac{(2-x^2)3x^2 - x^3(-2x)}{(2-x^2)^2} = \left(\frac{6x^2 - x^4}{(2-x^2)^2}\right)$	M1 A2,1,0	For quotient rule (or product rule on correct y)
(b)	$\frac{dy}{dx} = x \times \frac{1}{2}(4x+6)^{-0.5} \times 4 + (4x+6)^{0.5}$ $= \frac{6(x+1)}{(4x+6)^{0.5}} \quad \rightarrow \quad k = 6$	M1 A1 A1	product rule
4	$x(4-\sqrt{3}) = 13$ $x = \frac{13\left(4+\sqrt{3}\right)}{\left(4-\sqrt{3}\right)\left(4+\sqrt{3}\right)}$ $= 4 + \sqrt{3}$ $y = 1 - 2\sqrt{3}$	M1 A1 M1 A1 A1	eliminate y or x simplified rationalisation


© Cambridge International Examinations 2015

5	$(x-3)(x-3)(x-1)=0$	M1	
	$x^3-7x^2+15x-9=0$		
	$a=-7$	A1	
	$b=15$	A1	
	$c=-9$	A1	**AG** for c
6	$\log_x 2=\dfrac{\log_2 2}{\log_2 x}$	B1	
	$2\log_2 x=\log_2 x^2$	B1	
	$3=\log_2 8$	B1	
	$8x^2-29x+15\ (=0)$	M1	obtain quadratic and attempt to solve
	$\rightarrow (8x-5)(x-3)\ (=0)$		
	$x=\dfrac{5}{8}$ or $x=3$	A1	
7 **(i)**	$a=-\dfrac{20}{(t+2)^3}$	M1 A1	$k(t+2)^{-3}$ oe $k=-20$
	$t=3 \rightarrow a=-0.16$ m/s^2	A1FT	
(ii)	$\dfrac{10}{(t+2)^2}$ is never zero.	B1	
(iii)	$s=-\dfrac{10}{t+2}+5$	M1	integrate $\dfrac{k}{t+2}$
		A1	$k=-10$
		A1	+5
(iv)	$s=\left[-\dfrac{10}{t+2}\right]_3^8=-1+2$	M1	insert limits and subtract
	$=1$	A1	

© Cambridge International Examinations 2015

8	**(i)**	$\sec^2 x + \text{cosec}^2 x = \dfrac{1}{\cos^2 x} + \dfrac{1}{\sin^2 x}$	B1	
		$= \dfrac{\sin^2 x + \cos^2 x}{\sin^2 x \cos^2 x}$	B1	add fractions
		$= \dfrac{1}{\sin^2 x \cos^2 x}$	B1	use of $\sin^2 x + \cos^2 x = 1$
		$= \sec^2 x \text{cosec}^2 x$	B1	fully correct solution
	(ii)	$\dfrac{1}{\cos^2 x \sin^2 x} = 4\dfrac{\sin^2 x}{\cos^2 x}$	M1	
		$\rightarrow \quad 4\sin^2 x = 1$	A1	correct simplified equation
		$\sin x = \pm\dfrac{1}{\sqrt{2}}$		
		$x = 135°, 225°$	A1, A1	
9	**(i)**	$\text{f}(x) = 3x^2 + 12x + 2 = 3(x+2)^2 - 10$		
		$a = 3$	B1	
		$b = 2$	B1	
		$c = -10$	B1	
	(ii)	minimum $\text{f}(x) = -10$	B1FT	
		at $x = -2$	B1FT	
	(iii)	$\text{f}\left(\dfrac{1}{y}\right) = 0 \quad \rightarrow \quad \left(\dfrac{1}{y}\right) = (\pm)\sqrt{\dfrac{10}{3}} - 2$	M1	obtain explicit expression for $\dfrac{1}{y}$ or y
		$y = -5.74, \quad -0.26$	A1, A1	

© Cambridge International Examinations 2015

10	**(i)**	$\frac{\mathrm{d}}{\mathrm{d}x}(\mathrm{e}^{2-x^2}) = -2x\mathrm{e}^{2-x^2}$	B1	$k = -2$
	(ii)	$-\frac{3\mathrm{e}^{2-x^2}}{2} + c$	M1 A1FT	$D\mathrm{e}^{2-x^2}$ $D = \frac{-3}{2}$ or $\frac{3}{k}$
	(iii)	$\left[-\frac{3\mathrm{e}^{2-x^2}}{2}\right]_1^{\sqrt{2}} = -\frac{3}{2} + \frac{3}{2}\mathrm{e}$ 2.58	M1 A1	insert limits on *their* (ii) and subtract
	(iv)	$y = 3x\mathrm{e}^{2-x^2}$ $\frac{\mathrm{d}y}{\mathrm{d}x} = 3x(-2x\mathrm{e}^{2-x^2}) + 3\mathrm{e}^{2-x^2}$ $\frac{\mathrm{d}y}{\mathrm{d}x} = 0 \rightarrow x = \pm\frac{1}{\sqrt{2}} = \pm 0.707$ $y = \pm\frac{3}{\sqrt{2}}\mathrm{e}^{1.5} = \pm 9.51$	M1 A1 A1 A1	product rule both x or a pair both y
11	**(i)**	$\log N = \log A - t\log b$	B1	
	(ii)	t: 1 2 3 4 5 6 $\log N$: 3.30 3.11 2.95 2.77 2.60 2.41 $\ln N$: 7.60 7.17 6.79 6.38 5.98 5.56 	M1 M1 A1	find logs of N plot $\log N$ or $\ln N$ against t or $-t$ straight line passing through five points
	(iii)	gradient $= -\log b = \frac{2.415 - 3.3}{5} \rightarrow b = 1.5$ intercept $= \log A = 3.47 \rightarrow A = 2950$	DM1 DM1 A1	set gradient $= -\log b$ and solve set intercept $= \log A$ and solve both values correct
	(iv)	$t = 10 \rightarrow N = \frac{2950}{1.5^{10}} = 51$	B1	
	(v)	$N = 10 \rightarrow 1.5^t = 295 \rightarrow t = \frac{\log 295}{\log 1.5}$ $= 14$ years	M1 A1	substitute $N = 10$, *their* A, b into given or transformed equation

© Cambridge International Examinations 2015

12	$v_p = \begin{pmatrix} 250\cos 20^\circ \\ 250\sin 20^\circ \end{pmatrix}$, $v_r = \begin{pmatrix} V\cos 30^\circ \\ V\sin 30^\circ \end{pmatrix}$, $v_w = \begin{pmatrix} 0 \\ w \end{pmatrix}$	B1	
	$v_r = v_p + v_w$ $\begin{pmatrix} V\cos 30^\circ \\ V\sin 30^\circ \end{pmatrix} = \begin{pmatrix} 250\cos 20^\circ \\ 250\sin 20^\circ \end{pmatrix} + \begin{pmatrix} 0 \\ w \end{pmatrix}$		
	$V = \dfrac{250\cos 20^\circ}{\cos 30^\circ}$ $= 271\text{km/hr}$	M1 A1	equate x components and solve
	$w = V\sin 30^\circ - 250\sin 20^\circ$ $= 50.1\text{km/hr}$	M1 A1	equate y components and solve
	OR triangle with sides 250 V w opposite angles 60° 110° 10°	B1	
	sine rule: $\dfrac{w}{\sin 10^\circ} = \dfrac{250}{\sin 60^\circ}$ $w = 50.1\text{km/hr}$	M1 A1	apply to correct triangle and solve
	$\dfrac{V}{\sin 110^\circ} = \dfrac{250}{\sin 60^\circ}$ $V = 271\text{km/hr}$	M1 A1	apply to correct triangle and solve

© Cambridge International Examinations 2015