CAMBRIDGE INTERNATIONAL EXAMINATIONS

International General Certificate of Secondary Education

MARK SCHEME for the May/June 2014 series

0607 CAMBRIDGE INTERNATIONAL MATHEMATICS

0607/23 Paper 2 (Extended), maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2014 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

CAMBRIDGE
International Examinations

1		5.6[0]	**2**	**M1** for 7×0.8 oe
2	**(a)**	6.3×10^{-3}	**1**	
	(b)	5.94	**2**	**B1** for 0.24×10^9 or 57×10^8 or figures 594 seen
3	**(a)**	23	**1**	
	(b)	35	**1**	
	(c)	972	**1**	
4	**(a)**	$36x^2 - 4y^2$ oe	**2**	**B1** for $36x^2$ or $4y^2$
	(b)	$\sqrt{\frac{A+4y^2}{36}}$ or $\frac{1}{6}\sqrt{\mathrm{A}+(2\mathrm{y})^2}$ oe	**3**	**M1** for correct division by 36 or 6 **M1** for correct re-arrangement **M1** for correct square root
5	**(a)**	1	**1**	
	(b)	$3y^9$	**2**	**B1** for ky^9 or $3y^k$, $k \neq 0$
6	**(a)**	[x=] 32 [y=] 32	**1** **1FT**	**FT** *their x*
	(b)	[v=] 25 [w=] 65	**1** **1**	**FT** (90 – *their v*)
7		90	**3**	**B2** for $\frac{5}{8}\times x^2$ oe or $\frac{y}{40}=\left(\frac{12}{8}\right)^2$ or **M1** for $y = kx^2$, $k \neq 0, 1$
8	**(a)**	$10-\sqrt{2}$	**3**	**B2** for $12 - 4\sqrt{2} + 3\sqrt{2} - 2$ or better or **B1** if three of these terms correct
	(b)	$2\sqrt{5}$	**2**	**M1** for $\times\frac{\sqrt{5}}{\sqrt{5}}$ or $\frac{2\times5}{\sqrt{5}}$
9	**(a)**	Good sketch of stretch factor 2, x-axis invariant	**1**	
	(b)	Good sketch of translation $\begin{pmatrix}-1\\0\end{pmatrix}$	**1**	

© Cambridge International Examinations 2014

10		$\frac{2\sqrt{2}}{3}$ or $\frac{\sqrt{8}}{3}$	**3**	**B2** for $\sqrt{3^2 - 1^2}$ or better or **B1** for $k^2 + 1^2 = 3^2$ or better
11	**(a)**	$\cup$	**1**	
	(b)	$\cap$	**1**	
	(c)	$\in$ or $\{e\} \subset$	**1**	
	(d)	$\cup$	**1**	
12	**(a)**	3	**2**	**B1** for [f(1) =]4
	(b)	$\frac{12}{x}$	**1**	

© Cambridge International Examinations 2014