UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
International General Certificate of Secondary Education

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER			

2845008432

CHEMISTRY **0620/31**

Paper 3 (Extended) **May/June 2013**

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use a pencil for any diagrams, graphs or rough working.
Do not use staples, paper clips, highlighters, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.
Electronic calculators may be used.
A copy of the Periodic Table is printed on page 12.
You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.

This document consists of **12** printed pages.

© UCLES 2013

For Examiner's Use

1 Petroleum contains hydrocarbons which are separated by fractional distillation.

(a) (i) Complete the following definition of a hydrocarbon.

A hydrocarbon is a compound which .....

..... [2]

(ii) Explain what is meant by the term *fractional distillation*.

.....

.....

..... [2]

(b) Some of the fractions obtained from petroleum are given below.
State a use for each fraction.

bitumen .....

lubricating fraction .....

paraffin fraction .....

gasoline fraction ..... [4]

[Total: 8]

2 An element, **M**, has the electron distribution 2 + 8 + 18 + 3.

(a) Which group in the Periodic Table is element **M** likely to be in?

..... [1]

(b) Predict whether element **M** is a poor or a good conductor of electricity.
Give a reason for your answer.

..... [1]

(c) Binary compounds contain two atoms per molecule, for example HC*l*.
Identify an element which could form a binary compound with element **M**.

..... [1]

(d) Predict the formula of the sulfate of **M**. The formula of the sulfate ion is SO_4^{2-}.

..... [1]

© UCLES 2013

For Examiner's Use

(e) The hydroxide of **M** is a white powder which is insoluble in water.
Describe how you could show that this hydroxide is amphoteric.

.....

.....

..... [2]

[Total: 6]

3 A small piece of marble, $CaCO_3$, was added to 5.0 cm^3 of hydrochloric acid, concentration 1.0 mol / dm^3, at 25 °C. The time taken for the reaction to stop was measured. The experiment was repeated using 5.0 cm^3 of different solutions of acids. The acid was in excess in all of the experiments.

Typical results are given in the table.

experiment	temperature / °C	acid solution	time / min
1	25	hydrochloric acid 1.0 mol / dm^3	3
2	25	hydrochloric acid 0.5 mol / dm^3	7
3	25	ethanoic acid 1.0 mol / dm^3	10
4	15	hydrochloric acid 1.0 mol / dm^3	8

(a) (i) Explain why it is important that the pieces of marble are the same size and the same shape.

.....

.....

..... [2]

(ii) How would you know when the reaction had stopped?

..... [1]

(b) The equation for the reaction in experiment 1 is:

$$CaCO_3(s) + 2HCl(aq) \rightarrow CaCl_2(aq) + CO_2(g) + H_2O(l)$$

Complete the following ionic equation.

$$CaCO_3(s) + 2H^+(aq) \rightarrow \text{.....} + \text{.....} + \text{.....}$$

[1]

© UCLES 2013

[Turn over

(c) (i) Explain why the reaction in experiment 1 is faster than the reaction in experiment 2.

.....

..... [1]

(ii) The acids used for experiment 1 and experiment 3 have the same concentration. Explain why experiment 3 is slower than experiment 1.

.....

.....

..... [2]

(iii) Explain in terms of collisions between reacting particles why experiment 4 is slower than experiment 1.


.....

.....

..... [3]

[Total: 10]

4 The structural formula of cyclohexane is drawn below.



(a) The name gives information about the structure of the compound.
Hex because there are six carbon atoms and **cyclo** because they are joined in a ring.
What information about the structure of this compound is given by the ending **ane**?

.....

..... [2]

(b) What are the molecular and empirical formulae of cyclohexane?

molecular formula .....

empirical formula ..... [2]

© UCLES 2013

For Examiner's Use

(c) Draw the structural formula of cyclobutane.

[1]

(d) **(i)** Deduce the molecular formula of hexene.

..... [1]

(ii) Explain why cyclohexane and the alkene, hexene, are isomers.

.....

.....

..... [2]

(e) Describe a test which would distinguish between cyclohexane and the unsaturated hydrocarbon hexene.

test .....

result of test with cyclohexane .....

.....

result of test with hexene .....

..... [3]

[Total: 11]

© UCLES 2013

[Turn over

For Examiner's Use

5 The reactivity series shows the metals in order of reactivity.

(a) The reactivity series can be established using displacement reactions. A piece of zinc is added to aqueous lead nitrate. The zinc becomes coated with a black deposit of lead.

$$Zn + Pb^{2+} \rightarrow Zn^{2+} + Pb$$

Zinc is more reactive than lead.

The reactivity series can be written as a list of ionic equations.

.....	$\rightarrow$	.....	+	.....	most reactive metal : the best reductant (reducing agent)
Zn	$\rightarrow$	Zn^{2+}	+	$2e^-$	
Fe	$\rightarrow$	Fe^{2+}	+	$2e^-$	
Pb	$\rightarrow$	Pb^{2+}	+	$2e^-$	
Cu	$\rightarrow$	Cu^{2+}	+	$2e^-$	
Ag	$\rightarrow$	Ag^+	+	e^-	

(i) In the space at the top of the list, write an ionic equation for a metal which is more reactive than zinc. [1]

(ii) Write an ionic equation for the reaction between aqueous silver(I) nitrate and zinc.

..... [2]

(iii) Explain why the positive ions are likely to be oxidants (oxidising agents).

..... [1]

(iv) Deduce which ion is the best oxidant (oxidising agent).

..... [1]


(v) Which ion(s) in the list can oxidise lead metal?

..... [1]

© UCLES 2013

For Examiner's Use

(b) A reactivity series can also be established by measuring the voltage of simple cells. The diagram shows a simple cell.



Results from cells using the metals tin, cadmium, zinc and copper are given in the table below.

cell	electrode 1 positive electrode	electrode 2 negative electrode	voltage / volts
1	copper	cadmium	0.74
2	copper	tin	0.48
3	copper	zinc	1.10

Write the four metals in order of increasing reactivity and explain how you used the data in the table to determine this order.

.....

.....

..... [3]

[Total: 9]

6 Ammonia is a compound which only contains the elements nitrogen and hydrogen. It is a weak base.

(a) (i) Define the term *base*.

..... [1]

(ii) Given aqueous solutions of ammonia and sodium hydroxide, both having a concentration of 0.1 mol / dm^3, how could you show that ammonia is the weaker base?

.....

.....

..... [2]

For Examiner's Use

(b) Ammonia is manufactured by the Haber Process. The economics of this process require that as much ammonia as possible is made as quickly as possible.
Explain how this can be done using the following information.

The conditions for the following reversible reaction are:

- 450 °C
- 200 atmospheres pressure
- iron catalyst

$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ the reaction is exothermic

.....

.....

.....

.....

.....


..... [5]

(c) Another compound which contains only nitrogen and hydrogen is hydrazine, N_2H_4.

Complete the equation for the preparation of hydrazine from ammonia.

$$.....NH_3 + NaC\mathit{l}O \rightarrow N_2H_4 + ... + H_2O$$ [2]

(d) The structural formula of hydrazine is given below.



Draw a diagram showing the arrangement of the valency electrons in one molecule of the covalent compound hydrazine.
Use x to represent an electron from a nitrogen atom.
Use o to represent an electron from a hydrogen atom.

[3]

© UCLES 2013

(e) Hydrazine is a weak base and it removes dissolved oxygen from water. It is added to water in steel boilers to prevent rusting.

(i) One way it reduces the rate of rusting is by changing the pH of water.
What effect would hydrazine have on the pH of water?

..... [1]


(ii) Give a reason, other than pH, why hydrazine reduces the rate of rusting.

..... [1]

[Total: 15]

7 The hydroxides of the Group I metals are soluble in water. Most other metal hydroxides are insoluble in water.

(a) (i) Crystals of lithium chloride can be prepared from lithium hydroxide by titration.



25.0 cm^3 of aqueous lithium hydroxide is pipetted into the conical flask.
A few drops of an indicator are added. Dilute hydrochloric acid is added slowly to the alkali until the indicator just changes colour. The volume of acid needed to neutralise the lithium hydroxide is noted.

A neutral solution of lithium chloride, which still contains the indicator, is left. Describe how you could obtain a neutral solution of lithium chloride which does **not** contain an indicator.

.....

..... [2]

© UCLES 2013

(ii) You cannot prepare a neutral solution of magnesium chloride by the same method. Describe how you could prepare a neutral solution of magnesium chloride.

.....

.....

..... [3]

(b) The concentration of the hydrochloric acid was 2.20 mol / dm^3. The volume of acid needed to neutralise the 25.0 cm^3 of lithium hydroxide was 20.0 cm^3. Calculate the concentration of the aqueous lithium hydroxide.

$$LiOH + HCl \rightarrow LiCl + H_2O$$

.....

.....

..... [2]

(c) Lithium chloride forms three hydrates. They are $LiCl.H_2O$, $LiCl.2H_2O$ and $LiCl.3H_2O$. Which **one** of these three hydrates contains 45.9 % of water?
Show how you arrived at your answer.

.....

.....

..... [3]

[Total: 10]

8 There are three types of giant structure - ionic, metallic and giant covalent.

(a) In an ionic compound, the ions are held in a lattice by strong forces.

(i) Explain the term *lattice*.

.....

..... [2]

(ii) Explain how the ions are held together by strong forces.

.....

..... [1]

© UCLES 2013

(b) Describe the bonding in a typical metal.

.....

.....

..... [3]

(c) The electrical conductivities of the three types of giant structure are given in the following table.

type of structure	conductivity of solid	conductivity of liquid
ionic	poor	good
metallic	good	good
giant covalent	poor	poor

Explain the differences in electrical conductivity between the three types of giant structure and the difference, if any, between the solid and liquid states of the same structure.

.....

.....

.....

.....

..... [5]

[Total: 11]

© UCLES 2013

DATA SHEET
The Periodic Table of the Elements

	Group																	
	I	II									III	IV	V	VI	VII	0
																4 **He** Helium 2
				1 **H** Hydrogen 1							11 **B** Boron 5	12 **C** Carbon 6	14 **N** Nitrogen 7	16 **O** Oxygen 8	19 **F** Fluorine 9	20 **Ne** Neon 10
	7 **Li** Lithium 3	9 **Be** Beryllium 4									27 **Al** Aluminium 13	28 **Si** Silicon 14	31 **P** Phosphorus 15	32 **S** Sulfur 16	35.5 **Cl** Chlorine 17	40 **Ar** Argon 18
	23 **Na** Sodium 11	24 **Mg** Magnesium 12									70 **Ga** Gallium 31	73 **Ge** Germanium 32	75 **As** Arsenic 33	79 **Se** Selenium 34	80 **Br** Bromine 35	84 **Kr** Krypton 36
	39 **K** Potassium 19	40 **Ca** Calcium 20	45 **Sc** Scandium 21	48 **Ti** Titanium 22	51 **V** Vanadium 23	52 **Cr** Chromium 24	55 **Mn** Manganese 25	56 **Fe** Iron 26	59 **Co** Cobalt 27	59 **Ni** Nickel 28	64 **Cu** Copper 29	65 **Zn** Zinc 30	112 **Cd** Cadmium 48	119 **Sn** Tin 50	122 **Sb** Antimony 51	128 **Te** Tellurium 52	127 **I** Iodine 53	131 **Xe** Xenon 54
	85 **Rb** Rubidium 37	88 **Sr** Strontium 38	89 **Y** Yttrium 39	91 **Zr** Zirconium 40	93 **Nb** Niobium 41	96 **Mo** Molybdenum 42	**Tc** Technetium 43	101 **Ru** Ruthenium 44	103 **Rh** Rhodium 45	106 **Pd** Palladium 46	108 **Ag** Silver 47	112 **Cd** Cadmium 48	115 **In** Indium 49	119 **Sn** Tin 50	122 **Sb** Antimony 51	128 **Te** Tellurium 52	127 **I** Iodine 53	131 **Xe** Xenon 54
	133 **Cs** Caesium 55	137 **Ba** Barium 56	139 **La** Lanthanum 57 *	178 **Hf** Hafnium 72	181 **Ta** Tantalum 73	184 **W** Tungsten 74	186 **Re** Rhenium 75	190 **Os** Osmium 76	192 **Ir** Iridium 77	195 **Pt** Platinum 78	197 **Au** Gold 79	201 **Hg** Mercury 80	204 **Tl** Thallium 81	207 **Pb** Lead 82	209 **Bi** Bismuth 83	**Po** Polonium 84	**At** Astatine 85	**Rn** Radon 86
	Fr Francium 87	226 **Ra** Radium 88	227 **Ac** Actinium 89 †															

140 **Ce** Cerium 58	141 **Pr** Praseodymium 59	144 **Nd** Neodymium 60	**Pm** Promethium 61	150 **Sm** Samarium 62	152 **Eu** Europium 63	157 **Gd** Gadolinium 64	159 **Tb** Terbium 65	162 **Dy** Dysprosium 66	165 **Ho** Holmium 67	167 **Er** Erbium 68	169 **Tm** Thulium 69	173 **Yb** Ytterbium 70	175 **Lu** Lutetium 71
232 **Th** Thorium 90	**Pa** Protactinium 91	238 **U** Uranium 92	**Np** Neptunium 93	**Pu** Plutonium 94	**Am** Americium 95	**Cm** Curium 96	**Bk** Berkelium 97	**Cf** Californium 98	**Es** Einsteinium 99	**Fm** Fermium 100	**Md** Mendelevium 101	**No** Nobelium 102	**Lr** Lawrencium 103

*58-71 Lanthanoid series
†90-103 Actinoid series

Key: a = relative atomic mass; X = atomic symbol; b = proton (atomic) number

The volume of one mole of any gas is 24 dm^3 at room temperature and pressure (r.t.p.).

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© UCLES 2013