UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education Ordinary Level

CANDIDATE NAME

CENTRE NUMBER

CANDIDATE NUMBER

ADDITIONAL MATHEMATICS **4037/12**

Paper 1 **May/June 2012**

2 hours

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use a pencil for any diagrams or graphs.
Do not use staples, paper clips, highlighters, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 80.

For Examiner's Use	
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
Total	

This document consists of **20** printed pages.

© UCLES 2012

UNIVERSITY of CAMBRIDGE International Examinations

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1}\, b + \binom{n}{2} a^{n-2}\, b^2 + \ldots + \binom{n}{r} a^{n-r}\, b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} bc \sin A$$

© UCLES 2012

1 **(i)** Find $\int \sqrt{7x-5}\,\mathrm{d}x$. [3]

For Examiner's Use

(ii) Hence evaluate $\int_2^3 \sqrt{7x-5}\,\mathrm{d}x$. [2]

2 Using the substitution $u = 2^x$, find the values of x such that $2^{2x+2} = 5\,(2^x) - 1$. [5]

For Examiner's Use

3 Show that $\cot A + \dfrac{\sin A}{1 + \cos A} = \operatorname{cosec} A$. [4]

For Examiner's Use

4 Solve the simultaneous equations $5x + 3y = 2$ and $\frac{2}{x} - \frac{3}{y} = 1$. [5]

For Examiner's Use

5 Differentiate the following with respect to x.

For Examiner's Use

(i) $(2 - x^2)\ln(3x + 1)$ [3]

(ii) $\dfrac{4 - \tan 2x}{5x}$ [3]

6 You must not use a calculator in this question.

For Examiner's Use

(i) Express $\frac{8}{\sqrt{3}+1}$ in the form $a\left(\sqrt{3}-1\right)$, where a is an integer. [2]

An equilateral triangle has sides of length $\frac{8}{\sqrt{3}+1}$.

(ii) Show that the height of the triangle is $6-2\sqrt{3}$. [2]

(iii) Hence, or otherwise, find the area of the triangle in the form $p\sqrt{3} - q$, where p and q are integers. [2]

For Examiner's Use


7 **(i)** Sketch the graph of $y = |x^2 - x - 6|$, showing the coordinates of the points where the curve meets the coordinate axes. [3]

For Examiner's Use

(ii) Solve $|x^2 - x - 6| = 6$. [3]

8

For Examiner's Use



The figure shows a circle, centre O, with radius 10 cm. The lines XA and XB are tangents to the circle at A and B respectively, and angle AOB is $\frac{2\pi}{3}$ radians.

(i) Find the perimeter of the shaded region. [3]

(ii) Find the area of the shaded region. [4]

9 Variables N and x are such that $N = 200 + 50e^{\frac{x}{100}}$.

For Examiner's Use

(i) Find the value of N when $x = 0$. [1]

(ii) Find the value of x when $N = 600$. [3]

© UCLES 2012

(iii) Find the value of N when $\frac{dN}{dx} = 45$. [4]

For Examiner's Use

10 **(a)** It is given that $f(x) = \frac{1}{2+x}$ for $x \neq -2$, $x \in \mathbb{R}$.

For Examiner's Use

(i) Find $f''(x)$. [2]

(ii) Find $f^{-1}(x)$. [2]

(iii) Solve $f^2(x) = -1$. [3]

(b) The functions g, h and k are defined, for $x \in \mathbb{R}$, by

$$g(x) = \frac{1}{x+5}, \quad x \neq -5,$$
$$h(x) = x^2 - 1,$$
$$k(x) = 2x + 1.$$

Express the following in terms of g, h and/or k.

For Examiner's Use

(i) $\dfrac{1}{(x^2-1)+5}$ [1]

(ii) $\dfrac{2}{x+5} + 1$ [1]

11 The point P lies on the line joining $A(-1, -5)$ and $B(11, 13)$ such that $AP = \frac{1}{3}AB$.

For Examiner's Use

(i) Find the equation of the line perpendicular to AB and passing through P. [5]

The line perpendicular to AB passing through P and the line parallel to the x-axis passing through B intersect at the point Q.

(ii) Find the coordinates of the point Q. [2]

© UCLES 2012

(iii) Find the area of the triangle PBQ. [2]

For Examiner's Use

Answer only **one** of the following two alternatives.

For Examiner's Use

12 EITHER

At 12 00 hours, a ship has position vector $(54\mathbf{i} + 16\mathbf{j})$ km relative to a lighthouse, where **i** is a unit vector due East and **j** is a unit vector due North. The ship is travelling with a speed of 20 km h^{-1} in the direction $3\mathbf{i} + 4\mathbf{j}$.

(i) Show that the position vector of the ship at 15 00 hours is $(90\mathbf{i} + 64\mathbf{j})$ km. [2]

(ii) Find the position vector of the ship t hours after 12 00 hours. [2]


A speedboat leaves the lighthouse at 14 00 hours and travels in a straight line to intercept the ship. Given that the speedboat intercepts the ship at 16 00 hours, find

(iii) the speed of the speedboat, [3]

(iv) the velocity of the speedboat relative to the ship, [1]

(v) the angle the direction of the speedboat makes with North. [2]

OR



The position vectors of points A and B relative to an origin O are **a** and **b** respectively. The point P is such that $\overrightarrow{OP} = \frac{5}{4}\overrightarrow{OB}$. The point Q is such that $\overrightarrow{AQ} = \frac{1}{3}\overrightarrow{AB}$. The point R lies on OA such that RQP is a straight line where $\overrightarrow{OR} = \lambda\overrightarrow{OA}$ and $\overrightarrow{QR} = \mu\overrightarrow{PR}$.

(i) Express $\overrightarrow{OQ}$ and $\overrightarrow{PQ}$ in terms of **a** and **b**. [2]

(ii) Express $\overrightarrow{QR}$ in terms of λ, **a** and **b**. [2]

(iii) Express $\overrightarrow{QR}$ in terms of μ, **a** and **b**. [3]

(iv) Hence find the value of λ and of μ. [3]

© UCLES 2012

Start your answer to Question 12 here.

Indicate which question you are answering.

EITHER	
OR	

For Examiner's Use

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Continue your answer here if necessary.

For Examiner's Use

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© UCLES 2012