UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education Ordinary Level

CANDIDATE NAME

CENTRE NUMBER | | | | | |

CANDIDATE NUMBER | | | | |

9722452658

ADDITIONAL MATHEMATICS **4037/21**

Paper 2 **May/June 2013**

2 hours

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use a pencil for any diagrams or graphs.
Do not use staples, paper clips, highlighters, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 80.

This document consists of **16** printed pages.

© UCLES 2013

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Binomial Theorem

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \ldots + \binom{n}{r} a^{n-r} b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$.

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} bc \sin A$$


© UCLES 2013

1 Prove that $\left(\frac{1+\sin\theta}{\cos\theta}\right)^2 + \left(\frac{1-\sin\theta}{\cos\theta}\right)^2 = 2 + 4\tan^2\theta$. [4]

For Examiner's Use

2

For Examiner's Use



The velocity-time graph represents the motion of a particle moving in a straight line.

(i) Find the acceleration during the first 5 seconds. [1]

(ii) Find the length of time for which the particle is travelling with constant velocity. [1]

(iii) Find the total distance travelled by the particle. [3]


© UCLES 2013

3 Variables x and y are related by the equation $y = 10 - 4\sin^2 x$, where $0 \leqslant x \leqslant \frac{\pi}{2}$. Given that x is increasing at a rate of 0.2 radians per second, find the corresponding rate of change of y when $y = 8$. [6]

For Examiner's Use

4 **(i)** Sketch the graph of $y = |4x - 2|$ on the axes below, showing the coordinates of the points where the graph meets the axes. [3]

For Examiner's Use



(ii) Solve the equation $|4x - 2| = x$. [3]

5

For Examiner's Use



A piece of wire of length 96 cm is formed into the rectangular shape $PQRSTU$ shown in the diagram. It is given that $PQ = TU = SR = x$ cm. It may be assumed that PQ and TU coincide and that TS and QR have the same length.

(i) Show that the area, A cm^2, enclosed by the wire is given by $A = \dfrac{96x - 3x^2}{2}$. [2]

(ii) Given that x can vary, find the stationary value of A and determine the nature of this stationary value. [4]

6 Find the equation of the normal to the curve $y = \frac{x^2 + 8}{x - 2}$ at the point on the curve where $x = 4$. [6]

For Examiner's Use

7 **(i)** Find the first four terms in the expansion of $(2 + x)^6$ in ascending powers of x. [3]

For Examiner's Use

(ii) Hence find the coefficient of x^3 in the expansion of $(1 + 3x)(1 - x)(2 + x)^6$. [4]

8 The line $y = 2x - 8$ cuts the curve $2x^2 + y^2 - 5xy + 32 = 0$ at the points A and B. Find the length of the line AB. [7]

For Examiner's Use

© UCLES 2013

9 It is given that $x \in \mathbb{R}$ and that $\mathscr{E} = \{x : -5 < x < 12\}$,
$S = \{x : 5x + 24 > x^2\}$,
$T = \{x : 2x + 7 > 15\}$.

For Examiner's Use

Find the values of x such that

(i) $x \in S$, [3]

(ii) $x \in S \cup T$, [2]

(iii) $x \in (S \cap T)'$. [3]

© UCLES 2013

10 A plane, whose speed in still air is $240\,\text{kmh}^{-1}$, flies directly from A to B, where B is 500 km from A on a bearing of 032°. There is a constant wind of $50\,\text{kmh}^{-1}$ blowing from the west.

For Examiner's Use

(i) Find the bearing on which the plane is steered. [4]

© UCLES 2013

(ii) Find, to the nearest minute, the time taken for the flight. [4]

For Examiner's Use

11 A one-one function f is defined by $f(x) = (x - 1)^2 - 5$ for $x \geqslant k$.

For Examiner's Use

(i) State the least value that k can take. [1]


For this least value of k

(ii) write down the range of f, [1]

(iii) find $f^{-1}(x)$, [2]

(iv) sketch and label, on the axes below, the graph of $y = \mathrm{f}(x)$ and of $y = \mathrm{f}^{-1}(x)$, [2]

For Examiner's Use



(v) find the value of x for which $\mathrm{f}(x) = \mathrm{f}^{-1}(x)$. [2]

Question 12 is printed on the next page.

12 The function $f(x) = x^3 + x^2 + ax + b$ is divisible by $x - 3$ and leaves a remainder of 20 when divided by $x + 1$.

For Examiner's Use

(i) Show that $b = 6$ and find the value of a. [4]

(ii) Using your value of a and taking b as 6, find the non-integer roots of the equation $f(x) = 0$ in the form $p \pm \sqrt{q}$, where p and q are integers. [5]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© UCLES 2013