**CAMBRIDGE INTERNATIONAL EXAMINATIONS**

**GCE Ordinary Level**

## MARK SCHEME for the May/June 2014 series

# 4037 ADDITIONAL MATHEMATICS

**4037/11** Paper 1, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.


Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2014 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

CAMBRIDGE
International Examinations

| | | | |
|---|---|---|---|
| **1** | $\text{LHS} = \dfrac{\sin\ \theta}{\cos\ \theta} + \dfrac{\cos\ \theta}{1+\sin\ \theta}$ | **B1** | **B1** for use of $\tan\theta = \dfrac{\sin\theta}{\cos\theta}$ |
| | $= \dfrac{\sin\theta(1+\sin\theta) + \cos^2\theta}{\cos\theta(1+\sin\theta)}$ | **M1** | **M1** for attempt to obtain a single fraction |
| | $= \dfrac{1+\sin\theta}{\cos\theta(1+\sin\theta)}$ | **DM1** | **DM1** for use of $\sin^2\theta + \cos^2\theta = 1$ |
| | $= \dfrac{1}{\cos\theta}$ leading to $\sec\theta$ | **A1** | **A1** for 'finishing off' |
| | **Alternative solution**: $\text{LHS} = \dfrac{\sin\theta}{\cos\theta} + \dfrac{\cos\theta\,(1-\sin\theta)}{(1+\sin\theta)(1-\sin\theta)}$ | **B1** | **B1** for use of $\tan\theta = \dfrac{\sin\theta}{\cos\theta}$ |
| | $= \dfrac{\sin\theta}{\cos\theta} + \dfrac{\cos\theta(1-\sin\theta)}{\cos^2\theta}$ | **M1** | **M1** for multiplication by $(1-\sin\theta)$ |
| | $= \dfrac{\sin\theta}{\cos\theta} + \dfrac{(1-\sin\theta)}{\cos\theta}$ | **DM1** | **DM1** for use of $\sin^2\theta + \cos^2\theta = 1$ |
| | $= \dfrac{1}{\cos\theta}$ leading to $\sec\theta$ | **A1** | **A1** for 'finishing off' |
| | **Alternative solution**: $\text{LHS} = \dfrac{\tan\theta(1+\sin\theta) + \cos\theta}{1+\sin\theta}$ | **M1** | **M1** for attempt to obtain a single fraction |
| | $= \dfrac{\dfrac{\sin\theta}{\cos\theta} + \dfrac{\sin^2}{\cos\theta} + \cos\theta}{1+\sin\theta}$ | **B1** | **B1** for use of $\tan\theta = \dfrac{\sin\theta}{\cos\theta}$ |
| | $= \dfrac{\sin\theta + \sin^2\theta + \cos^2\theta}{\cos\theta(1+\sin\theta)}$ | | |
| | $= \dfrac{1+\sin\theta}{\cos\theta(1+\sin\theta)}$ | **DM1** | **DM1** for use of $\sin^2\theta + \cos^2\theta = 1$ |
| | $= \dfrac{1}{\cos\theta}$ leading to $\sec\theta$ | **A1** | **A1** for 'finishing off' |



© Cambridge International Examinations 2014

| | | | | |
|---|---|---|---|---|
| **2** | **(i)** | $\lvert\mathbf{a}\rvert = \sqrt{4^2 + 3^2} = 5$<br>$\lvert\mathbf{b} + \mathbf{c}\rvert = \sqrt{(-3)^2 + 4^2} = 5$ | **M1**<br><br>**A1** | **M1** for finding the modulus of either **a** or **b** + **c**<br><br>**A1** for completion |
| | **(ii)** | $\lambda\begin{pmatrix}4\\3\end{pmatrix} + \mu\begin{pmatrix}2\\2\end{pmatrix} = 7\begin{pmatrix}-5\\2\end{pmatrix}$<br><br>$4\lambda + 2\mu = -35$ and $3\lambda + 2\mu = 14$<br><br>leading to $\lambda = -49$, $\mu = 80.5$ | <br><br>**M1**<br><br>**DM1**<br><br>**A1** | <br><br>**M1** for equating like vectors and obtaining 2 linear equations<br><br>**DM1** for solution of simultaneous equations<br>**A1** for both |
| **3** | **(a)** | **(i)** **(ii)** **(iii)**<br> | **B1**<br>**B1**<br>**B1** | **B1** for each |
| | **(b) (i)** | 2 | **B1** | |
| | **(ii)** | 0 | **B1** | |
| **4** | | $k(4x - 3) = 4x^2 + 8x - 8$<br>$4x^2 + x(8 - 4k) + 3k - 8 = 0$<br>$b^2 - 4ac = (8 - 4k)^2 - 16(3k - 8)$<br>$= 16k^2 - 112k + 192$<br>$b^2 - 4ac < 0$, $k^2 - 7k + 12 < 0$<br><br>critical values $k = 3, 4$<br><br>$\therefore 3 < k < 4$ | **M1**<br><br>**DM1**<br><br>**DM1**<br><br>**A1**<br><br>**A1** | **M1** for equating the line and the curve and attempt to obtain a quadratic equation in $k$<br><br>**DM1** for use of $b^2 - 4ac$ with $k$<br><br>**DM1** for solution of a 3 term quadratic equation, dependent on both previous M marks<br>**A1** for both critical values<br><br>**A1** for the range |
| **5** | **(i)** | $\frac{dy}{dx} = 2xe^{x^2}$ | **B1B1** | **B1** for $e^{x^2}$, **B1** for $2xe^{x^2}$ |
| | **(ii)** | $\frac{1}{2}e^{x^2}$ | **M1A1** | **M1** for $ke^{x^2}$ **A1** for $\frac{1}{2}e^{x^2}$ |
| | **(iii)** | $\left(\frac{1}{2}e^4\right) - \left(\frac{1}{2}\right) = 26.8$ | **DM1**<br>**A1** | **DM1** for correct use of limits<br>**A1** for 26.8, allow exact value |

© Cambridge International Examinations 2014

| | | | | |
|---|---|---|---|---|
| **6** | **(i)** | $\mathbf{AB} = \begin{pmatrix} 10 & 19 \\ 32 & 37 \\ 14 & 14 \end{pmatrix}$ | **M1**<br><br>**A1** | **M1** for at least 3 correct elements of a $3 \times 2$ matrix<br>**A1** for all correct |
| | **(ii)** | $\mathbf{B}^{-1} = \frac{1}{7}\begin{pmatrix} 5 & -1 \\ -3 & 2 \end{pmatrix}$ | **B1**<br>**B1** | **B1** for $\frac{1}{7}$, **B1** for $\begin{pmatrix} 5 & -1 \\ -3 & 2 \end{pmatrix}$ |
| | **(iii)** | $2\begin{pmatrix} 2 & 1 \\ 3 & 5 \end{pmatrix}\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -3 \\ -22 \end{pmatrix}$ | **M1** | **M1** for obtaining in matrix form |
| | | $\begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{7}\begin{pmatrix} 5 & -1 \\ -3 & 2 \end{pmatrix}\begin{pmatrix} -1.5 \\ -11 \end{pmatrix} = \frac{1}{7}\begin{pmatrix} 3.5 \\ -17.5 \end{pmatrix}$ | **M1** | **M1** for pre-multiplying by $\mathbf{B}^{-1}$ |
| | | $x = 0.5, y = -2.5$ | **A1** | **A1** for both |
| **7** | **(i)** | $y = 2x^2 - \frac{1}{x+1}(+c)$ | **B1**<br>**B1** | **B1** for each correct term |
| | | when $x = \frac{1}{2}, y = \frac{5}{6}$ so $\frac{5}{6} = \frac{1}{2} - \frac{2}{3} + c$ | **M1** | **M1** for attempt to find $+c$, must have at least 1 of the previous B marks |
| | | leading to $c = 1$ | **A1** | Allow **A1** for $c = 1$ |
| | | $\left(y = 2x^2 - \frac{1}{x+1} + 1\right)$ | | |
| | **(ii)** | When $x = 1, y = \frac{5}{2}$ | **M1** | **M1** for using $x = 1$ in their **(i)** to find $y$ |
| | | $\frac{dy}{dx} = \frac{17}{4}$ so gradient of normal $= -\frac{4}{17}$ | **B1** | **B1** for gradient of normal |
| | | Equation of normal $y - \frac{5}{2} = -\frac{4}{17}(x-1)$ | **DM1** | **DM1** for attempt at **normal** equation |
| | | $(8x + 34y - 93 = 0)$ | **A1** | **A1** – allow unsimplified<br>( fractions must not contain decimals) |

© Cambridge International Examinations 2014

| | | | | |
|---|---|---|---|---|
| **8** | **(i)** | $\log p = n\log V + \log k$<br><br>$\ln V$: 2.30, 3.91, 4.61, 5.30<br>$\ln p$: 4.55, 2.14, 1.10, 0.10<br><br>$\lg V$: 1, 1.70, 2, 2.30<br>$\lg p$: 1.98, 0.93, 0.48, 0.04<br><br><br> | **B1**<br><br><br><br><br><br><br><br><br><br>**M1**<br>**A2,1,0** | **B1** for statement, but may be implied by later work.<br><br><br><br><br><br><br><br><br><br>**M1** for plotting a suitable graph<br>–1 for each error in points plotted |
| | **(ii)** | Use of gradient = $n$<br>$n = -1.5$ (allow –1.4 to –1.6) | **DM1**<br>**A1** | **DM1** for equating numerical gradient to $n$ |
| | **(iii)** | Allow 13 to 16 | **DM1**<br>**A1** | **DM1** for use of *their* graph or substitution into *their* equation. |
| **9** | **(a)** | Distance travelled = area under graph<br>$= \frac{1}{2}(60+20)\times 12 = 480$ | **M1**<br><br>**A1** | **M1** for realising that area represents distance travelled and attempt to find area |
| | **(b)** | <br> | **B1**<br>**B1**<br><br>**B1** | **B1** for velocity of 2 $ms^{-1}$ for $0 \leqslant t \leqslant 6$<br>**B1** for velocity of zero for *their* '6' to *their* '25'<br>**B1** for velocity of 1 $ms^{-1}$ for $25 \leqslant t \leqslant 30$ |
| | **(c) (i)** | $v = 4 - \frac{16}{t+1}$<br><br>When $v = 0,\ t = 3$ | **M1**<br><br>**DM1**<br>**A1** | **M1** for attempt at differentiation<br><br>**DM1** for equating velocity to zero and attempt to solve |
| | **(ii)** | $a = \frac{16}{(t+1)^2}$<br><br>$0.25(t+1)^2 = 16$<br><br>$t = 7$ | **M1**<br><br><br><br>**A1** | **M1** for attempt at differentiation and equating to 0.25 with attempt to solve |

© Cambridge International Examinations 2014

| | | | |
|---|---|---|---|
| **10 (a)** | 1 digit even numbers 2 | **B1** | |
| | 2 digit even numbers $4 \times 2 = 8$ | **B1** | |
| | 3 digit even numbers $3 \times 3 \times 2 = 18$ | **B1** | |
| | Total = 28 | **B1** | |
| **(b) (i)** | 3M 5W = 35<br>4M 4W = 175<br>5M 3W = 210 | **B1**<br>**B1**<br>**B1** | |
| | Total = 420 | **B1** | **B1** for addition to obtain final answer, must be evaluated. |
| | **or** ${}^{12}C_8$ – 6M 2W – 7M 1W<br>495 – 70 – 5 = 420 | | **or**: as above, final **B1** for subtraction to get final answer |
| **(ii)** | Oldest man in, oldest woman out and vice–versa | | |
| | ${}^{10}C_7 \times 2 = 240$ | **B1, B1** | **B1** for ${}^{10}C_7$ , B1 for realising there are 2 identical cases |
| | **Alternative:**<br>1 man out 1 woman in<br>6 men 4 women | | |
| | 6M 1W : ${}^6C_6 \times {}^4C_1 = 4$<br>5M 2W : ${}^6C_5 \times {}^4C_2 = 36$<br>4M 3W : ${}^6C_4 \times {}^4C_3 = 60$<br>3M 4W : ${}^6C_3 \times {}^4C_4 = 20$ | | |
| | Total = 120 | **B1** | All separate cases correct for **B1** |
| | There are 2 identical cases to consider, so 240 ways in all. | **B1** | **B1** for realising there are 2 identical cases, which have integer values |

© Cambridge International Examinations 2014

| | | | |
|---|---|---|---|
| **11 (a)** | | | **In each case the last A mark is for a second correct solution and no extra solutions within the range** |
| | $5\sin 2x + 3\cos 2x = 0$ | **M1** | **M1** for use of tan |
| | $\tan 2x = -0.6$ | **DM1** | **DM1** for dealing with $2x$ correctly |
| | $2x = 149°, 329°$ | | |
| | $x = 74.5°, 164.5°$ | **A1,A1** | **A1** for each |
| | **Alternatives:** | | |
| | $\sin(2x + 31°) = 0$ or $\cos(2x - 59°) = 0$ | **M1** | **M1** for either, then mark as above |
| **(b)** | $2\cot^2 y + 3\operatorname{cosec} y = 0$ | | |
| | $2(\operatorname{cosec}^2 y - 1) + 3\operatorname{cosec} y = 0$ | **M1** | **M1** for use of correct identity |
| | $2\operatorname{cosec}^2 y + 3\operatorname{cosec} y - 2 = 0$ | | |
| | $(2\operatorname{cosec} y - 1)(\operatorname{cosec} y + 2) = 0$ | **M1** | **M1** for attempt to factorise a 3 term quadratic equation |
| | One valid solution | | |
| | $\operatorname{cosec} y = -2,\ \sin y = -\frac{1}{2}$ | | |
| | $y = 210°, 330°$ | **A1,A1** | **A1** for each |
| | **Alternative:** | | |
| | $2\frac{\cos^2 y}{\sin^2 y} + \frac{3}{\sin y} = 0$ | **M1** | **M1** for use of $\cot y = \frac{\cos y}{\sin y}$ and $\operatorname{cosec} y = \frac{1}{\sin y}$ |
| | leads to $2\sin^2 y - 3\sin y - 2 = 0$ | | |
| | and $\sin y = -\frac{1}{2}$ only | **M1** | **M1** for attempt to factorise a 3 term quadratic equation |
| | $y = 210°, 330°$ | **A1A1** | |
| **(c)** | $3\cos(z + 1.2) = 2$ | | |
| | $\cos(z + 1.2) = \frac{2}{3}$ | | |
| | $(z + 1.2) = 0.8411, 5.442, 7.124$ | **M1** | **M1** for correct order of operations to end up with 0.8411 radians or better |
| | $z = 4.24, 5.92$ | **A1**<br>**A1A1** | **A1** for one of 5.441 or 7.124 (or better)<br>**A1** for each valid solution |

© Cambridge International Examinations 2014