UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Ordinary Level

MARK SCHEME for the October/November 2011 question paper for the guidance of teachers

4037 ADDITIONAL MATHEMATICS

4037/22 Paper 2, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

UNIVERSITY of CAMBRIDGE
International Examinations

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Accuracy mark for a correct result or statement independent of method marks.

- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.

- The symbol √ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.

- Note: B2 or A2 means that the candidate can earn 2 or 0.
  B2, 1, 0 means that the candidate can earn anything from 0 to 2.

© University of Cambridge International Examinations 2011

The following abbreviations may be used in a mark scheme or used on the scripts:

AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)

BOD Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)

CAO Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)

ISW Ignore Subsequent Working

MR Misread

PA Premature Approximation (resulting in basically correct work that is insufficiently accurate)

SOS See Other Solution (the candidate makes a better attempt at the same question)

Penalties

MR –1 A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through √ " marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy.

OW –1,2 This is deducted from A or B marks when essential working is omitted.


PA –1 This is deducted from A or B marks in the case of premature approximation.

S –1 Occasionally used for persistent slackness – usually discussed at a meeting.

EX –1 Applied to A or B marks when extra solutions are offered to a particular equation. Again, this is usually discussed at the meeting.


© University of Cambridge International Examinations 2011

1 **(a)**



6 correctly positioned	B1
14 correctly positioned	B1
1 correctly positioned	B1
9 correctly positioned	B1√


(b)



B1

[5]

2 **(i)**



Sine curve through (0, 0), (90, 1)(180, 0)	B1
Cosine curve through (0, 2), (45, 1), (90, 0), (135, 1), (180, 2)	M1 A1
(ii) 2	B1√
(iii) 2√	B1√

[5]

3

$f\left(\frac{1}{2}\right) = 0$ or f(–2) = 25 (or f(–1) found)	M1
$\frac{1}{2} + \frac{1}{4}a - \frac{11}{2} + b = 0$ or $\frac{1}{4}a + b = 5$ or $a + 4b = 20$	A1
$-32 + 4a + 22 + b = 25$ or $4a + b = 35$	A1
Solve simultaneous equations	M1
$a = 8, b = 3$	A1
f(–1) = 18	A1√

[6]

© University of Cambridge International Examinations 2011

4	**(i)**	Matrix multiplication $\begin{pmatrix} 9 & -22 & 6 \\ 16 & -29 & 2 \end{pmatrix}$	M1 A1
	(ii)	Matrix multiplication $\begin{pmatrix} -18 \\ -2 \end{pmatrix}$	M1 A1
	(iii)	$-\frac{1}{13}\begin{pmatrix} -5 & 2 \\ -1 & 3 \end{pmatrix}$ or $\frac{1}{13}\begin{pmatrix} 5 & -2 \\ 1 & -3 \end{pmatrix}$	B1+B1 **[6]**
5	**(i)**	Evidence of $4 \times 3 \times 3 \times 2 \times 2(\times 1 \times 1)$ or $4 \times 3 \times 3 \times 2 \times 2(\times 1 \times 1)$ or $4! \times 3!$ 144	M1 A1
	(ii)	4! (or 24) for boys and 3! (or 6) for girls 288	M1 A1
	(iii)	Evidence of $4 \times (120) \times 3$ or $(4) \times 5! \times (3)$ 1440	M1 A1 **[6]**
6	**(i)**	$4x - 20 > 140$ oe	B1
	(ii)	$x(x - 10) < 3000$	B1
	(iii)	$x > 40$ Factorise 3 term quadratic –50 and 60 $40 < x < 60$	B1 M1 A1 A1 **[6]**
7	**(i)**	$\frac{1}{2}(14^2 - 6^2)\theta = 32$ 0.4	M1A1 A1
	(ii)	$\frac{CF}{2} = 20 \sin 0.2$ or $CF^2 = 20^2 + 20^2 - 2 \times 20 \times 20 \cos 0.4$ (CF=)7.95 Uses $s = r\theta$ Complete plan including $s = r\theta$ 25.5 or 25.6	M1 A1 M1 M1 A1 **[8]**

© University of Cambridge International Examinations 2011

8	**(i)**	Solve $\cos\left(\frac{t}{3}\right) = \frac{2}{12}$	M1
		4.21 or 4.2(0)	A1
	(ii)	$a = \frac{\mathrm{d}v}{\mathrm{d}t} = k\sin\frac{t}{3}$	M1
		$k = -4$	A1
		–3.36 or –3.37	A1√
	(iii)	Find t when $v = 0$	M1
		$s = \int v\mathrm{d}t = 36\sin\frac{t}{3}$	B1
		substitute t into $k\sin\frac{t}{3}$	M1
		36	A1√ **[9]**
9	**(i)**	$a = 2$, $b = -3$, $c = -8$ or $2(x-3)^2 - 8$	B1+B1+B1
	(ii)	graph with 3 distinct parts all above and touching x axis	B1
		quadratic shape for middle section reflected in x axis	B1
		cusp	B1
		completely correct	B1
	(iii)	identifies 0 and 8	M1
		$0 < k < 8$	A1 **[9]**
10	**(i)**	$3x^2 - 18x + 24$	B1
		Solves 3 term quadratic equals zero	M1
		$x = 4$ (and $x = 2$)	A1 cso
		$Q(4, 18)$	A1
	(ii)	$\frac{x^4}{4} - 3x^3 + 12x^2 + 2x$	B2, 1, 0
		Uses limits of 0 and x_Q in integral (area under curve = 72)	M1
		Area under PQ = (32 + 8)	B1
		Plan for area including use of limits	M1
		32	A1 **[10]**

© University of Cambridge International Examinations 2011

11E (i)	$\overrightarrow{OX} = \mathbf{a} + \mu(\mathbf{b}\text{-}\mathbf{a})$ or $(1-\mu)\mathbf{a} + \mu\mathbf{b}$	M1A1
(ii)	$\overrightarrow{OS} = \frac{3}{5}\mathbf{a}$	B1
	$\overrightarrow{OT} = \frac{7}{5}\mathbf{b}$	B1
	$\overrightarrow{OX} = \frac{3}{5}\mathbf{a} + \lambda\left(\frac{7}{5}\mathbf{b} - \frac{3}{5}\mathbf{a}\right)$ or $\overrightarrow{OX} = (1-\lambda)\frac{3}{5}\mathbf{a} + \lambda\frac{7}{5}\mathbf{b}$	M1A1
(iii)	Equate components or arrange to $(\alpha)\ \mathbf{a} = (\beta)\ \mathbf{b}$ and put $\alpha = \beta = 0$	M1
	Solve simultaneous equations	DM1
	$\lambda = \frac{1}{2}$	A1
	$\mu = \frac{7}{10}$	A1
		[10]
11O (i)	$\overrightarrow{OP} = 3\mathbf{c}$	B1
	$\overrightarrow{OQ} = \frac{3}{2}\mathbf{d}$	B1
	$\overrightarrow{DR} = \overrightarrow{CD} = \mathbf{d} - \mathbf{c}$	B1
	$\overrightarrow{OR} = \overrightarrow{OD} + \overrightarrow{DR} = 2\mathbf{d} - \mathbf{c}$	M1A1
(ii)	Finds **two** of $\overrightarrow{PQ},\ \overrightarrow{QR},\ \overrightarrow{PR}$	M1
	Two of $\overrightarrow{PQ} = \frac{3}{2}\mathbf{d} - 3\mathbf{c},\ \overrightarrow{QR} = \frac{1}{2}\mathbf{d} - \mathbf{c},\ \overrightarrow{PR} = 2\mathbf{d} - 4\mathbf{c}$	A1+A1
	Express one vector as multiple of another	M1
	3:1	A1
		[10]

© University of Cambridge International Examinations 2011