UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education Ordinary Level

CANDIDATE NAME

CENTRE NUMBER

CANDIDATE NUMBER

4611981584

ADDITIONAL MATHEMATICS **4037/13**

Paper 1 **October/November 2013**

2 hours

Candidates answer on the Question Paper.

No additional materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use a pencil for any diagrams or graphs.
Do not use staples, paper clips, highlighters, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 80.

DC (SLM) 81101
© UCLES 2013

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Binomial Theorem

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \ldots + \binom{n}{r} a^{n-r} b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$.

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} bc \sin A$$

© UCLES 2013

1 The coefficient of x^2 in the expansion of $(2 + px)^6$ is 60.

For Examiner's Use

(i) Find the value of the positive constant p. [3]

(ii) Using your value of p, find the coefficient of x^2 in the expansion of $(3 - x)(2 + px)^6$. [3]

© UCLES 2013

2 Solve $2\lg y - \lg(5y + 60) = 1$. [5]

For Examiner's Use

3 Show that $\tan^2\theta - \sin^2\theta = \sin^4\theta\sec^2\theta$. [4]

For Examiner's Use

For Examiner's Use

4 A curve has equation $y = \dfrac{e^{2x}}{(x+3)^2}$.

(i) Show that $\dfrac{dy}{dx} = \dfrac{Ae^{2x}(x+2)}{(x+3)^3}$, where A is a constant to be found. [4]

(ii) Find the exact coordinates of the point on the curve where $\dfrac{dy}{dx} = 0$. [2]

© UCLES 2013

5 For $x \in \mathbb{R}$, the functions f and g are defined by

$$f(x) = 2x^3,$$

$$g(x) = 4x - 5x^2.$$

For Examiner's Use


(i) Express $f^2\left(\frac{1}{2}\right)$ as a power of 2. [2]

(ii) Find the values of x for which f and g are increasing at the same rate with respect to x. [4]

6 Do not use a calculator in this question.

For Examiner's Use

The diagram shows part of the curve $y = 4 - x^2$.



Show that the area of the shaded region can be written in the form $\frac{\sqrt{2}}{p}$, where p is an integer to be found. [6]

7 It is given that $\mathbf{A} = \begin{pmatrix} 2t & 2 \\ t^2 - t + 1 & t \end{pmatrix}$.

For Examiner's Use

(i) Find the value of t for which $\det \mathbf{A} = 1$. [3]


(ii) In the case when $t = 3$, find $\mathbf{A}^{-1}$ and hence solve

$$3x + y = 5,$$
$$7x + 3y = 11.$$

[5]

8 The diagram shows two concentric circles, centre O, radii 4 cm and 6 cm. The points A and B lie on the larger circle and the points C and D lie on the smaller circle such that ODA and OCB are straight lines.

For Examiner's Use



(i) Given that the area of triangle OCD is 7.5 cm^2, show that $\theta = 1.215$ radians, to 3 decimal places. [2]

(ii) Find the perimeter of the shaded region. [4]

© UCLES 2013

(iii) Find the area of the shaded region. [3]

For Examiner's Use

9 **(a)** **(i)** Solve $6\sin^2 x = 5 + \cos x$ for $0° < x < 180°$. [4]

For Examiner's Use

(ii) Hence, or otherwise, solve $6\cos^2 y = 5 + \sin y$ for $0° < y < 180°$. [3]

© UCLES 2013

(b) Solve $4\cot^2 z - 3\cot z = 0$ for $0 < z < \pi$ radians. [4]

For Examiner's Use


10 The variables s and t are related by the equation $t = ks^n$, where k and n are constants. The table below shows values of variables s and t.

For Examiner's Use

s	2	4	6	8
t	25.00	6.25	2.78	1.56

(i) A straight line graph is to be drawn for this information with $\lg t$ plotted on the vertical axis. State the variable which must be plotted on the horizontal axis. [1]

(ii) Draw this straight line graph on the grid below. [3]



© UCLES 2013

(iii) Use your graph to find the value of k and of n. [4]

For Examiner's Use

(iv) Estimate the value of s when $t = 4$. [2]

Question 11 is printed on the next page.

© UCLES 2013

11 **(i)** Given that $\int_0^k \left(2e^{2x} - \frac{5}{2}e^{-2x}\right) dx = \frac{3}{4}$, where k is a constant, show that

$$4e^{4k} - 12e^{2k} + 5 = 0.$$ [5]

For Examiner's Use

(ii) Using a substitution of $y = e^{2k}$, or otherwise, find the possible values of k. [4]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© UCLES 2013