CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge Ordinary Level

MARK SCHEME for the October/November 2014 series

4037 ADDITIONAL MATHEMATICS

4037/22 Paper 2, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.



Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2014 series for most Cambridge IGCSE®, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

CAMBRIDGE
International Examinations

1 **(a)**		**B1**	
		**B1**	
(b)	No.in H only $= 50 - x$; No in F only $= 60 - x$ Sum: $50 - x + 60 - x + x + 30 - 2x = 98$ $x = 14$	**B1** **M1** **A1**	Both written or on diagram Add at least 3 terms each with x involved and equate to 98 soi
2	$9x^2 + 2x - 1 < (x+1)^2$ $8x^2 < 2$ oe isw $-\frac{1}{2} < x < \frac{1}{2}$	**M1** **A1** **A1**	Expand and collect terms
3	$\log_2(x+3) = \log_2 y + 2 \rightarrow x + 3 = 4y$ $\log_2(x+y) = 3 \rightarrow x + y = 8$ $x + 3 = 4(8 - x)$ $5x = 29 \rightarrow x = 5.8,$ oe $y = 2.2$ oe	**B1** **B1** **M1** **A1** **A1**	Eliminate y or x from two linear three term equations

© Cambridge International Examinations 2014

4	**(i)**	$\mathrm{f}(37)=3$ or $\mathrm{gf}(x)=\dfrac{\sqrt{x-1}-3-2}{2\left(\sqrt{x-1}-3\right)-3}$	**B1**	
		$\mathrm{gf}(37)=\dfrac{3-2}{6-3}=\dfrac{1}{3}$	**B1**	
	(ii)	$y=\sqrt{x-1}-3 \rightarrow (y+3)^2=x-1$	**M1**	Rearrange and square in any order
		$(x+3)^2+1=\mathrm{f}^{-1}(x)$ oe isw	**A1**	Interchange x and y and complete
	(iii)	$y=\dfrac{x-2}{2x-3}$		
		$2xy-3y=x-2 \rightarrow 2xy-x=3y-2$	**M1**	Multiply and collect like terms
		$\dfrac{3x-2}{2x-1}=\mathrm{g}^{-1}(x)$ oe	**A1**	Interchange and complete Mark final answer
5	**(i)**	$B=900$	**B1**	
	(ii)	$B=500+400\mathrm{e}^2=3455$ or 3456 or 3460	**B1**	3455.6 scores **B0**
	(iii)	$\left(\dfrac{\mathrm{d}B}{\mathrm{d}t}=\right)80\mathrm{e}^{0.2t}$	**B1**	
		$t=10 \rightarrow \dfrac{\mathrm{d}B}{\mathrm{d}t}=80\mathrm{e}^2=591\,(/\mathrm{day})$	**B1**	awrt
	(iv)	$10\,000=500+400\mathrm{e}^{0.2t} \rightarrow \mathrm{e}^{0.2t}=(23.75)$	**M1**	$\mathrm{e}^{0.2t}=k$
		$0.2t=\ln 23.75$	**DM1**	take logs: $0.2t=\ln k$
		$t=15.8\,(\mathrm{days})$	**A1**	awrt

© Cambridge International Examinations 2014

6 **(i)**	$(x+2)^2 + x^2 = 10$	**B1**	
	$x^2 + 2x - 3 = 0 \rightarrow (x+3)(x-1) = 0$	**M1**	3 term quadratic with attempt to solve
	Points (1, 3), (–3, –1) isw	**A1**	both x or a pair
		A1	both y or second pair
	or elimination of x leads to $y^2 - 2y - 3 = 0$, then as above		
(ii)	$m^2x^2 + 10mx + 25 + x^2 = 10$	**B1**	
	$(m^2+1)x^2 + 10mx + 15 = 0$		
	$b^2 - 4ac = (0) \rightarrow 100m^2 - 60(m^2+1) = 0$	**M1**	attempt to use discriminant on three term quadratic. Allow unsimplified
		A1	
	$m = \pm\sqrt{\frac{3}{2}}$ oe isw	**A1**	cao ± is required
	Alternative solution:		
	$\frac{dy}{dx} = \frac{-x}{\sqrt{10-x^2}}$ or $\frac{dy}{dx} = -\frac{x}{y}$	**B1**	allow unsimplified
	Result:		
	$y^2 = x^2 + 5y$ after inserted in $y = mx + 5$		
	Attempt to solve with $x^2 + y^2 = 10$	**M1**	Eliminate x or y
	$y = 2, x = \pm\sqrt{6}$	**A1**	both
	$m = \pm\frac{3}{\sqrt{6}}$ oe	**A1**	
7 **(i)**	$v = 2\cos t + 1$	**B1**	mark final answer
(ii)	$2\cos t + 1 = 0$	**M1**	equate their v to zero (must be a differential) and attempt to solve to find an **angle**
	$t = \frac{2\pi}{3}$ or 2.09	**A1**	awrt
(iii)	$t = \frac{2\pi}{3} \rightarrow x = 2\sin\left(\frac{2\pi}{3}\right) + \frac{2\pi}{3} = 3.83\,\text{m}$	**B1**	awrt
	$a = -2\sin t$	**B1ft**	ft *their* v (2nd differential)
	$t = \frac{2\pi}{3}\ a = -\sqrt{3} = -\frac{1.73}{4}\,\text{ms}^{-2}$	**DB1ft**	ft using *their* **angle** t in correct a awrt
8 **(i)**	$\frac{dy}{dx} = \frac{(2+x^2)\times 2x - x^2 \times 2x}{(2+x^2)^2} = \frac{4x}{(2+x^2)^2}$	**M1**	apply quotient or product rule
		A1	unsimplified
	$k = 4$	**A1**	k=4 does not need to be specifically identified
(ii)	$\int \frac{x}{(2+x^2)^2}\,dx = \frac{1}{4} \times \frac{x^2}{2+x^2} + (c)$ isw	**B1**	$\frac{1}{\text{their } k} \times$ original function
		B1	

© Cambridge International Examinations 2014

9	$\left(a+3\sqrt{5}\right)^2 = a^2 + 3\sqrt{5}a + 3\sqrt{5}a + 45$ oe	**B1**	anywhere
	Equate: $a^2 + a + 45 = 51$ and $6a - b = 0$	**B1** **B1**	
	$(a+3)(a-2) = 0$	**M1**	Attempt to solve three term quadratic with integer coefficients obtained by equating coeffs
	$a = -3, 2$ $b = -18, 12$	**A1** **A1**	Both *a*s correct or one correct pair Both *b*s correct
10 (i)	$\sec x \operatorname{cosec} x = \dfrac{1}{\cos x \sin x}$	**B1**	anywhere
	$\cot x = \dfrac{\cos x}{\sin x}$	**B1**	anywhere
	LHS $= \dfrac{1-\cos^2 x}{\cos x \sin x}$ oe	**B1ft**	correct addition of *their* terms
	$= \dfrac{\sin^2 x}{\cos x \sin x} = \tan x$ AG	**B1**	use of identity and cancel
(ii)	$3\cot x - \cot x = \tan x \rightarrow 2\cot x = \tan x$	**M1**	equate and collect like terms, allow sign errors
	$\tan^2 x = 2$ oe	**A1**	
	x = 54.7, 125.3, 234.7, 305.3	**A1** **A1**	2 values only 2 more values. awrt
11 (i)	Area of sector $= \dfrac{1}{2} \times x^2 \times 0.8 \left(= 0.4x^2 \text{ cm}^2\right)$	**B1**	anywhere
	$SR = 5\sin 0.8 \left(= 3.59\right)$ or $OR = 5\cos 0.8 \left(= 3.48\right)$	**B1**	*SR* may be seen in stated $\dfrac{1}{2}ab\sin C$
	Area of triangle = $\dfrac{1}{2} 5\cos 0.8 \times 5\sin 0.8 = 6.247 \text{ cm}^2$	**M1**	insert correct terms into correct area formulae
	$0.08x^2 = 6.247$	**A1**	
	$x = 8.837$ cm AG	**A1**	
(ii)	$SQ = 8.84 - 5 \left(= 3.84 \text{ cm}\right)$		
	$PR = 8.84 - 5\cos 0.8 \left(= 5.35 \text{ or } 5.36 \text{ cm}\right)$	**B1**	two lengths from *SQ, PR, PQ* awrt
	$PQ = 8.84 \times 0.8 \left(= 7.07 \text{ cm}\right)$	**B1**	third length awrt
	Perimeter = 19.84 to 19.86 cm or rounded to 19.8 or 19.9	**B1**	sum
(iii)	Area $PQSR = 4 \times 6.247$	**M1**	
	$= 25 \text{ cm}^2$	**A1**	24.95 to 25

© Cambridge International Examinations 2014

12 **(i)**	$f(2) = 3(2^3) - 14(2^2) + 32 = 0$ Or complete long division	**B1**	
(ii)	$f(x) = (x-2)(3x^2 - 8x - 16)$	**M1** **A1**	$3x^2$ and 16 $8x$ and correct signs
	$f(x) = (x-2)(x-4)(3x+4)$	**M1** **A1**	Factorise three term quadratic
(iii)	$x = 2, 4$	**B1**	
(iv)	$\int 3x - 14 + \frac{32}{x^2}\,dx = 1.5x^2 - 14x - \frac{32}{x}(+c)$	**B1** **B1**	first 2 terms third term correct unsimplified
	Area $= \left[1.5x^2 - 14x - \frac{32}{x}\right]_2^4$	**M1**	Limits of 2 and 4 and subtract
	$= (-)\ 2$	**A1**	

© Cambridge International Examinations 2014