CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge Ordinary Level

MARK SCHEME for the October/November 2015 series

4037 ADDITIONAL MATHEMATICS

4037/22 Paper 2, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2015 series for most Cambridge IGCSE®, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

CAMBRIDGE
International Examinations

Abbreviations

awrt answers which round to
cao correct answer only
dep dependent
FT follow through after error
isw ignore subsequent working
nfww not from wrong working
oe or equivalent
rot rounded or truncated
SC Special Case
soi seen or implied
www without wrong working

1 **(i)**	$f(-2) = -32 - 16 + 30 + 18 = 0$	B1	All four evaluated terms must be seen. Allow if correct long division used
(ii)	$f(x) = (x+2)(4x^2 - 12x + 9)$	M1 A1	Coefficients 4 and 9 Coefficient –12
	$= (x+2)(2x-3)(2x-3)$	A1	All three factors together
	$f(x) = 0 \rightarrow x = -2,\ 1.5$ nfww	A1	Allow 1.5 mentioned just once
2 **(i)**	$(2-3x)^6 = 64 - 576x + 2160x^2$ isw	B1B1B1	
(ii)	$2160 - 2 \times 576 = 1008$	M1 A1	*their* final 2160 + 2 × *their* final –576
3 **(i)**	$\overrightarrow{AB} = \begin{pmatrix} -15 \\ 8 \end{pmatrix}$	B1	Allow $\overrightarrow{BA}$ May be implied by later work.
	$\lvert AB \rvert = \sqrt{15^2 + 8^2}$ $(= 17)$	M1	Use of Pythagoras on *their AB*
	Speed $= 17 \times 3 = 51$km/hr	A1	Must be exact
(ii)	$\overrightarrow{BC} = \begin{pmatrix} 16 \\ -30 \end{pmatrix}$	B1	Allow $\overrightarrow{CB}$
	$\lvert BC \rvert = \sqrt{16^2 + 30^2}$ $(= 34)$	M1	Use of Pythagoras on *their BC*
	Time taken $= \frac{34}{51} \times 60 = 40$ mins (or $\frac{2}{3}$ hrs)	A1	Allow answers which round to 40 to 2sf. Accept 0.66 or 0.67 hrs. Mark final answer.

© Cambridge International Examinations 2015

4	**(a)**	$2\mathbf{BA} = 2\begin{pmatrix} 1 & -2 & 4 \\ -2 & 3 & 0 \end{pmatrix}\begin{pmatrix} 2 & -1 \\ 3 & 5 \\ 7 & 4 \end{pmatrix}$ $= 2\begin{pmatrix} 24 & 5 \\ 5 & 17 \end{pmatrix} = \begin{pmatrix} 48 & 10 \\ 10 & 34 \end{pmatrix}$	B3,2,1,0	–1 each error in 2×2 result. Failure to multiply by 2 is one error
	(b) (i)	$\mathbf{C}^{-1} = \frac{1}{8}\begin{pmatrix} 6 & -2 \\ 1 & 1 \end{pmatrix}$ isw	B1 B1	$\frac{1}{8}$ Matrix
	(ii)	$\mathbf{I} - \mathbf{D} = \begin{pmatrix} -2 & 2 \\ -1 & -3 \end{pmatrix}$	B1	
		$\mathbf{X} = \mathbf{C}^{-1}(\mathbf{I} - \mathbf{D}) = \frac{1}{8}\begin{pmatrix} 6 & -2 \\ 1 & 1 \end{pmatrix}\begin{pmatrix} -2 & 2 \\ -1 & -3 \end{pmatrix}$	M1	Pre multiply *their* **I – D** with *their* $\mathbf{C}^{-1}$
		$= \frac{1}{8}\begin{pmatrix} -10 & 18 \\ -3 & -1 \end{pmatrix}$ isw	A1	
5	**(a)**	$2^{3(q-1)} \times 2^{2p+1} = 2^{14}$	B1	Correct powers of 2 allow unsimplified isw
		$3^{2(p-4)} \times 3^{q} = 3^{4}$	B1	Correct powers of 3 allow unsimplified isw
		Solve $3q + 2p = 16$ $q + 2p = 12$	M1	Attempt to solve *their* linear equations by eliminating one variable
		$p = 5, \quad q = 2$	A1	Both correct
	(b)	$(3x - 2)(x + 1)$	M1	LHS oe isw
		$= 50$	A1	50 from correct processing of $2 - \lg 2$
		$3x^2 + x - 52 = 0 \rightarrow (3x + 13)(x - 4)$	M1	Solution of *their* three term quadratic Roots must be obtained from correct quadratic
		$x = 4$	A1	
		$x = -\frac{13}{3}$ discarded	A1	

© Cambridge International Examinations 2015

6 (i)	$a = 3, \quad b = 2, \quad c = 4$	B1B1B1	
(ii)	$\frac{dy}{dx} = 8\cos 4x$ isw	M1 A1FT	$\pm k\cos cx$ and no other term in x $\quad c \neq 1$ $bc \times \cos cx$ and no other term
(iii)	$x = \frac{\pi}{2} \rightarrow \frac{dy}{dx} = 8\cos 2\pi = 8$	DM1	Find *their* correct numerical $\frac{dy}{dx}$
	Eqn: $\frac{y-3}{x-\frac{\pi}{2}} = -\frac{1}{8} \qquad \left(\rightarrow y = -\frac{1}{8}x + 3.20\right)$	M1	Find equation with *their* numerical normal gradient ie $\frac{-1}{\frac{dy}{dx}}$ and point $\left(\frac{\pi}{2},\ 3\right)$
		A1	All correct isw
7 (i)	$\frac{h}{8} = \frac{6-r}{6} \rightarrow h = \frac{4}{3}(6-r)$	M1 A1	Uses correct ratio. Cannot be implied
(ii)	$V = \pi r^2 h = \pi r^2 \times \frac{4}{3}(6-r)$ $= 8\pi r^2 - \frac{4}{3}\pi r^3$	B1	**AG** all steps must be seen Penalise missing brackets at any point in working
(iii)	$\frac{dV}{dr} = 16\pi r - 4\pi r^2$	M1 A1	Differentiate at least one power reduced by one
	$\frac{dV}{dr} = 0 \rightarrow r = 4$	M1 A1	Attempt to solve – must get $r = \ldots$ Correct value of r. Ignore $r = 0$
	$V = \frac{128}{3}\pi \qquad (= 42.7\pi)$	A1	Correct value of V. Condone 134.
	$\frac{d^2V}{dr^2} = 16\pi - 8\pi r < 0$ when $r = 4 \rightarrow$ max	B1	$\frac{d^2V}{dr^2}$ must be correct and some indication of a negative value seen plus maximum stated

© Cambridge International Examinations 2015

8	**(i)**	Gradient $AB = \frac{8-2}{9+3}$ $\left(=\frac{1}{2}\right)$ isw	B1	
		Equation AB and $x=0 \rightarrow \frac{y-2}{0+3}=\frac{1}{2}$ $\left(\rightarrow y=\frac{1}{2}x+3.5\right)$	M1	Find equation with *their* gradient and set $x = 0$
		$\rightarrow y = 3.5$	A1	
	(ii)	D is (3, 5)	B1	
	(iii)	Gradient perpendicular = –2	M1	Use of $m_1 \times m_2 = -1$ on gradient used for *their* line in **(i)**
		Equation perpendicular $\frac{y-5}{x-3}=-2$ $\rightarrow (y=-2x+11)$	A1	
	(iv)	E is (0, 11)	A1FT	
	(v)	Area of $ABE = \frac{1}{2}\begin{vmatrix} -3 & 9 & 0 & -3 \\ 2 & 8 & 11 & 2 \end{vmatrix}$	M1	For area of ABE or ECD. $\frac{1}{2}$ and *their* correct 8 elements must be seen.
		$=\frac{1}{2}\lvert -24+99-18+33 \rvert = 45$	A1	45 condone from E (0, –4)
		Area of $EDC = \frac{1}{2}\begin{vmatrix} 3 & 0 & 0 & 3 \\ 5 & 3.5 & 11 & 5 \end{vmatrix}$		
		$=\frac{1}{2}\lvert -10.5+33 \rvert = 11.25$	A1	11.25 condone from E(0, –4)

© Cambridge International Examinations 2015

9 (i)	$\tan 2x = -\frac{5}{4}$ $(2x = 128.7,\ 308.7)$	M1	For obtaining and using $\tan 2x = \pm\frac{5}{4}$ or $\pm\frac{4}{5}$ resulting in $2x =$
	$x = 64.3$ awrt 154.3 awrt	A1 A1FT	tan*x* = … gets M0 *their* $64.3° + 90°$
(ii)	$\text{cosec}^2 y + 3\text{cosec}y - 4 = 0$ or $4\sin^2 y - 3\sin y - 1 = 0$ $(\text{cosec}y + 4)(\text{cosec}y - 1) = 0$ or $(4\sin y + 1)(\sin y - 1) = 0$	B1	In any form as a three term quadratic.
	$\sin y = -\frac{1}{4}$ or $\sin y = 1$	M1	Solve three term quadratic in $\text{cosec}\, y$ or $\sin y$
	$y = 194.5,\ 345.5,\ 90$	A1A1A1	Answers must be obtained from the correct quadratic
(iii)	$z + \frac{\pi}{4} = \pi - \frac{\pi}{3}$ or	B1	Accept 2.09, 2.10, $\pi - 1.05$, $\pi - 1.04$ on RHS. Could be implied by final answer
	$z + \frac{\pi}{4} = \pi + \frac{\pi}{3}$	B1	Accept 4.19, 4.18, $\pi + 1.05$, $\pi + 1.04$ on RHS. Could be implied by final answer
	$z = \frac{5\pi}{12},\ \frac{13\pi}{12}$	B1B1	Answers must be correct multiples of π.
10 (i)	$s = \frac{1}{2}e^{2t} + 3e^{-2t} - t + (c)$	M1	Integrate : coefficient of $\frac{1}{2}$ or 3 seen with no change in powers of e. Ignore $-t$
	$t = 0,\ s = 0 \rightarrow c = -3.5$ $\left(s = \frac{1}{2}e^{2t} + 3e^{-2t} - t - 3.5\right)$	A1 A1	All correct and simplified
(ii)	$v = 0 \rightarrow u^2 - u - 6 = 0$ oe	M1	Obtain three term quadratic in u or e^{2t} Condone sign errors.
	$(u - 3)(u + 2) = 0$	DM1	Solve three term quadratic
	$\rightarrow u = 3 \rightarrow t = \frac{1}{2}\ln 3$ or 0.549	A1	Accept 0.55 No second answer
(iii)	$t = \frac{1}{2}\ln 3 \rightarrow a = 2e^{2t} + 12e^{-2t}$ $= 6 + 4 = 10$	B1 B1	Correct differentiation Allow awrt 10.0 or 9.99. No second answer.

© Cambridge International Examinations 2015