Cambridge International Examinations
Cambridge Ordinary Level

ADDITIONAL MATHEMATICS **4037/23**

Paper 2 **October/November 2016**

MARK SCHEME

Maximum Mark: 80

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2016 series for most Cambridge IGCSE®, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

© UCLES 2016

Abbreviations

awrt answers which round to
cao correct answer only
dep dependent
FT follow through after error
isw ignore subsequent working
oe or equivalent
rot rounded or truncated
SC Special Case
soi seen or implied
www without wrong working

Question	Answer	Mark	Part Marks
1	$\frac{\left(\sqrt{5}+3\sqrt{3}\right)}{\left(\sqrt{5}+\sqrt{3}\right)}\times\frac{\left(\sqrt{5}-\sqrt{3}\right)}{\left(\sqrt{5}-\sqrt{3}\right)}$	**M1**	rationalise with $\left(\sqrt{5}-\sqrt{3}\right)$
	$=\frac{5+3\sqrt{15}-\sqrt{15}-9}{5-3}$	**A1**	numerator (3 or 4 terms)
	$=\frac{2\sqrt{15}-4}{2}=\sqrt{15}-2$	**A1**	denominator and completion
2	$\ln e^{3x}=\ln 6e^{x}$		
	$3x=\ln 6e^{x}$	**M1**	one law of indices/logs
	$3x=\ln 6+\ln e^{x}$	**M1**	second law of indices/logs
	$3x=\ln 6+x$		
	$x=\frac{1}{2}\ln 6$ or $\ln\sqrt{6}$ or 0.896	**A1**	www oe in base 10
3 (i)	$\frac{\mathrm{d}}{\mathrm{d}x}\left(\frac{\sin x}{1+\cos x}\right)=\frac{(1+\cos x)\cos x+\sin x\sin x}{(1+\cos x)^2}$	**M1**	Quotient Rule (or Product Rule from $(\sin x)(1+\cos x)^{-1}$)
		A1	correct unsimplified
	$=\frac{\cos x+\cos^2 x+\sin^2 x}{(1+\cos x)^2}$	**B1**	use of $\sin^2 x+\cos^2 x=1$ oe
	$=\frac{1+\cos x}{(1+\cos x)^2}$	**A1**	completion
(ii)	$\int_0^2\left(\frac{1}{1+\cos x}\right)\mathrm{d}x=\left[\frac{\sin x}{1+\cos x}\right]_0^2$	**M1**	correct integrand
	awrt 1.56	**A1**	

© UCLES 2016

Question	Answer	Mark	Part Marks
4 (i)	$p(2)=0 \rightarrow 8+4a+2b-24=0$	**B1**	
	$\rightarrow (4a+2b=16)$		
	$p(1)=-20 \rightarrow 1+a+b-24=-20$	**B1**	
	$\rightarrow (a+b=3)$		
		M1	solve *their* linear equations for a or b
	$a=5$ and $b=-2$	**A1**	
(ii)	$p(x)=x^3+5x^2-2x-24$	**M1**	find quadratic factor
	$=(x-2)(x^2+7x+12)$	**A1**	correct quadratic factor soi
	$=(x-2)(x+3)(x+4)$	**M1**	factorise quadratic factor and write as product of 3 linear factors
	$p(x)=0 \rightarrow x=2, -3, -4.$	**A1**	if 0 scored, **SC2** for roots only
5 (i)	$AB^2=\left(\sqrt{3}+1\right)^2+\left(\sqrt{3}-1\right)^2$	**M1**	use cosine rule
	$-2\left(\sqrt{3}+1\right)\left(\sqrt{3}-1\right)\cos 60$		
	$=3+1+2\sqrt{3}+3+1-2\sqrt{3}-2$	**A1**	at least 7 terms
	$=6$	**A1**	correct completion AG
(ii)	$\frac{\sin A}{\sqrt{3}-1}=\frac{\sin 60}{\sqrt{6}}$	**M1**	sine rule (or cosine rule)
	$\sin A=\frac{\left(\sqrt{3}-1\right)\sin 60}{\sqrt{6}}=\frac{\sqrt{6}-\sqrt{2}}{4}$ oe or 0.259 or 0.2588…	**A1**	correct explicit expression for $\sin A$ AG
(iii)	Area $=\frac{1}{2}\left(\sqrt{3}+1\right)\left(\sqrt{3}-1\right)\sin 60$	**M1**	correct substitution into $\frac{1}{2}ab\sin C$
	$=\frac{\sqrt{3}}{2}$	**A1**	
6 (i)	$\frac{dy}{dx}=\sec^2 x$	**B1**	
	$x=\frac{\pi}{4} \rightarrow \frac{dy}{dx}=\sec^2\frac{\pi}{4}=2$	**B1**	evaluated
	$y=8$	**B1**	
	Equation of tangent $\frac{y-8}{x-\frac{\pi}{4}}=2$	**B1**	
	$(4-2y=\pi-16,\ y=2x+6.429\ldots,$ $\frac{\pi}{4}=0.7853\ldots)$		

© UCLES 2016

Question	Answer	Mark	Part Marks
(ii)	$\sec^2 x = \tan x + 7$ $\tan^2 x - \tan x - 6 = 0$ oe $(\tan x - 3)(\tan x + 2) = 0$ $\tan x = 3$ or $\tan x = -2$ $x = 1.25,\quad 2.03$	**M1** **M1** **A1A1**	use $\sec^2 x = 1 + \tan^2 x$ to obtain a 3 term quadratic in tan*x* solve three term quadratic for tan*x* extras in range lose final **A1**
7 (i)	$r^2 + h^2 = (0.5h + 2)^2$ oe $r^2 = 0.25h^2 + 2h + 4 - h^2$ $r^2 = 2h + 4 - 0.75h^2$	**M1** **A1**	correct expansion and r^2 subject and completion www AG
(ii)	$V = \frac{1}{3}\pi r^2 h = \frac{\pi}{3}(2h^2 + 4h - 0.75h^3)$ $\frac{dV}{dh} = \frac{\pi}{3}(4h + 4 - 2.25h^2)$ $\frac{dv}{dh} = 0 \rightarrow 2.25h^2 - 4h - 4 = 0$ $h = 2.49$ only	**B1** **M1** **A1** **M1** **A1**	any correct form in terms of *h* only differentiate *V* correct differentiation equate to 0 and solve 3 term quadratic cao
(iii)	$\frac{d^2V}{dh^2} = \frac{\pi}{3}(4 - 4.5h)$ when $h = 2.49$ $(-7.545\ldots) < 0$ so maximum	**M1** **A1**	differentiate *their* 3 term $\frac{dV}{dh}$ and substitute *their h* draw correct conclusion www
8 (i)	$\cos TOA = \frac{6}{10} \rightarrow$ $TOA = 0.927$	**M1** **A1**	any method
(ii)	area of major sector = $\frac{1}{2}6^2(2\pi - 2 \times their\, 0.927)$ (= 79.7) area of half kite = $\frac{1}{2}(6)\sqrt{10^2 - 6^2}$ (=24) area of kite × 2 (=48) complete correct plan awrt 128	**M2** **M1** **DM1** **DM1** **A1**	or **M1** for $\frac{1}{2}\ 6^2(2 \times their\ 0.927)$ **DM1** for $\pi \times 6^2 - \frac{1}{2}\ 6^2(2 \times their\ 0.927)$ any method *their* major sector + *their* kite
(iii)	arc length = $6 \times (2\pi - 2 \times their\, 0.927) + 2 \times \sqrt{10^2 - 6^2}$) awrt 42.6	**M1** **A1**	complete correct method

© UCLES 2016

Question	Answer	Mark	Part Marks
9 (i)	$p = 4$	**B1**	
(ii)	$\tan\alpha = \pm\frac{1}{3}$ or ±3 or 18.4° or 71.6° seen	**M1**	could use cos or sin
	108	**A1**	
(iii)	$\boldsymbol{r}_A = \begin{pmatrix}1\\5\end{pmatrix} + t\begin{pmatrix}their\ p\\-3\end{pmatrix}$	**B1**	
(iv)	$\boldsymbol{r}_B = \begin{pmatrix}q\\-15\end{pmatrix} + t\begin{pmatrix}3\\-1\end{pmatrix}$	**B1**	
(v)	$5 - 3t = -15 - t$	**M1**	$\boldsymbol{r}_A = \boldsymbol{r}_B$ and equate y/**j** and solve for t
	$\rightarrow t = 10$	**A1**	
(vi)	$\begin{pmatrix}41\\-25\end{pmatrix}$ only	**B1**	
(vii)	$q = 11$ only	**B1**	
10 (i)	$\text{fg}(x) = \ln\left(2\text{e}^x + 3\right) + 2$	**B1**	isw
(ii)	$\text{ff}(x) = \ln\left(\ln x + 2\right) + 2$	**B1**	isw
(iii)	$x = 2\text{e}^y + 3$	**M1**	change x and y and make e^y the subject
	$\text{e}^y = \frac{x-3}{2}$		
	$\text{g}^{-1}(x) = \ln\left(\frac{x-3}{2}\right)$ oe	**A1**	
(iv)	e^2 or 7.39	**B1**	
(v)	$\text{gf}(x) = 2\text{e}^{(\ln x+2)} + 3 = 20$	**B1**	gf correct and equation set up correctly
	$2\text{e}^{\ln x}\text{e}^2 + 3 = 20$	**M1**	one law of indices/logs
	$2x\text{e}^2 = 17$	**M1**	second law of indices/logs
	$x = \frac{17}{2\text{e}^2}$ or 1.15	**A1**	www if 0 scored, **SC2** for 17.3…

© UCLES 2016

Question	Answer	Mark	Part Marks
11 (i)	$\mathbf{A}^2 = \begin{pmatrix} 2 & q \\ p & 3 \end{pmatrix}\begin{pmatrix} 2 & q \\ p & 3 \end{pmatrix} = \begin{pmatrix} 4+pq & 2q+3q \\ 2p+3p & pq+9 \end{pmatrix}$	B2,1,0	–1 each error
	$\mathbf{A}^2 - 5\mathbf{A} = 2\mathbf{I} \rightarrow 4 + pq - 10 = 2$ or $9 + pq - 15 = 2$	M1	equate top left or bottom right elements
	$\rightarrow pq = 8$	A1	accept $p = \frac{8}{q}, \quad q = \frac{8}{p}$
(ii)	$\det \mathbf{A} = 6 - pq$	B1	
	$6 - pq = -3p$ and solve	M1	*their* $\det \mathbf{A} = -3p$ and use *their* $pq = k$ oe to solve for p or q
	$\rightarrow p = \frac{2}{3}$	A1	
	$q = 12$	A1	**FT** from *their* $pq = k$

© UCLES 2016